This is an electronic reprint of the original article.
This reprint may differ from the original in pagination and typographic detail.

Gabányi, K. É.; Frey, S.; Paragi, Z.; Järvelä, E.; Morokuma, T.; An, T.; Tanaka, M.; Tar, I.
The radio structure of the peculiar narrow-line Seyfert 1 galaxy candidate J1100+4421

Published in:
Monthly Notices of the Royal Astronomical Society

DOI:
[10.1093/mnras/stx2449](https://doi.org/10.1093/mnras/stx2449)

Published: 11/01/2018

Document Version
Publisher's PDF, also known as Version of record

Please cite the original version:
Gabányi, K. É., Frey, S., Paragi, Z., Järvelä, E., Morokuma, T., An, T., Tanaka, M., & Tar, I. (2018). The radio structure of the peculiar narrow-line Seyfert 1 galaxy candidate J1100+4421. *Monthly Notices of the Royal Astronomical Society*, 473(2), 1554-1561. <https://doi.org/10.1093/mnras/stx2449>

The radio structure of the peculiar narrow-line Seyfert 1 galaxy candidate J1100+4421

K. É. Gabányi,^{1,2★} S. Frey,^{1★} Z. Paragi,³ E. Järvelä,^{4,5} T. Morokuma,⁶ T. An,^{7,8} M. Tanaka⁹ and I. Tar¹⁰

¹Konkoly Observatory, MTA Research Centre for Astronomy and Earth Sciences, Konkoly Thege Miklós út 15-17, H-1121 Budapest, Hungary

²MTA-ELTE Extragalactic Astrophysics Research Group, ELTE TTK Pázmány Péter sétány 1/A, H-1117 Budapest, Hungary

³Joint Institute for VLBI ERIC, PO Box 2, NL-7990 AA Dwingeloo, the Netherlands

⁴Aalto University Metsähovi Radio Observatory, Metsähovintie 114, FI-02540 Kylmäla, Finland

⁵Aalto University Department of Electronics and Nanoengineering, PO Box 15500, FI-00076 Aalto, Finland

⁶Institute of Astronomy, Graduate School of Science, University of Tokyo, 2-21-1, Osawa, Mitaka, Tokyo 181-0015, Japan

⁷Shanghai Astronomical Observatory, Chinese Academy of Sciences, 80 Nandan Road, 200030 Shanghai, P.R. China

⁸Key Laboratory of Radio Astronomy, Chinese Academy of Sciences, 210008 Nanjing, P.R. China

⁹National Astronomical Observatory of Japan, Mitaka, Tokyo 181-8588, Japan

¹⁰Department of Astronomy, Eötvös University, PO Box 32, H-1518 Budapest, Hungary

Accepted 2017 September 19. Received 2017 September 19; in original form 2017 April 19

ABSTRACT

Narrow-line Seyfert 1 galaxies (NLS1) are an intriguing subclass of active galactic nuclei. Their observed properties indicate low central black hole mass and high accretion rate. The extremely radio-loud NLS1 sources often show relativistic beaming and are usually regarded as younger counterparts of blazars. Recently, the object SDSS J110006.07+442144.3 was reported as a candidate NLS1 source. The characteristics of its dramatic optical flare indicated its jet-related origin. The spectral energy distribution of the object was similar to that of the γ -ray detected radio-loud NLS1, PMN J0948+0022. Our high-resolution European very long baseline interferometry network observations at 1.7 and 5 GHz revealed a compact core feature with a brightness temperature of $\gtrsim 10^{10}$ K. Using the lowest brightness temperature value and assuming a moderate Lorentz factor of ~ 9 , the jet viewing angle is $\lesssim 26^\circ$. Archival Very Large Array data show a large-scale radio structure with a projected linear size of ~ 150 kpc reminiscent of double-sided morphology.

Key words: galaxies: active – galaxies: individual: SDSS J110006.07+442144.3 – galaxies: Seyfert.

1 INTRODUCTION

Narrow-line Seyfert 1 galaxies (NLS1, e.g. Pogge 2000) form a special subclass of active galactic nuclei (AGNs). They were first recognized as Seyfert 1 galaxies with unusually narrow H I lines (Osterbrock & Pogge 1985). NLS1 sources are identified by three features: (i) narrow permitted optical lines, with full width at half-maximum, $\text{FWHM}(\text{H}\beta) < 2000 \text{ km s}^{-1}$, (ii) a flux ratio of $[\text{O III}] \lambda 5007$ to $\text{H}\beta$ smaller than 3, and (iii) the strong emission feature caused by Fe II multiplets. The latter indicates a direct view to the accretion disc, although very recently, based upon the spectroscopic studies of a large sample of NLS1 candidate sources, Cracco et al. (2016) concluded that NLS1 sources do not necessarily have strong Fe II emission lines. The relatively narrow widths of the permitted lines are explained with the lower velocities of the

clouds in the broad-line region orbiting a low-mass black hole, 10^6 – $10^8 M_\odot$ (Mathur 2000). These low-mass black holes in NLS1 sources accrete at high rates, close to the Eddington limit (e.g. Collin & Kawaguchi 2004, and references therein).

Concerning the radio emission, statistical studies showed that only ~ 7 per cent of NLS1 sources are radio loud (e.g. Komossa et al. 2006; Zhou et al. 2006). The radio-loudness is commonly assessed following the original prescription of Kellermann et al. (1989), using the ratio of the 6-cm radio flux density to the $4400 \text{ \AA}$ optical flux density (R), and sources are usually classified to be radio-loud if $R \gtrsim 10$ (e.g. Komossa et al. 2006). Most of the radio-loud NLS1 (RLNLS1) sources have steep spectra and resemble the compact steep-spectrum sources (Komossa et al. 2006). However, the extremely radio-loud NLS1s (~ 2.5 per cent of NLS1 sources, with $R \geq 100$; Komossa et al. 2006) show blazar-like properties: flat radio spectrum, compact radio cores, substantial variability, high brightness temperatures, flat X-ray spectra and blazar-like spectral energy distribution (e.g. Yuan et al. 2008). Thus, these

* E-mail: krisztina.g@gmail.com (KÉG); frey.sandor@csfk.mta.hu (SF)

objects are thought to possess relativistic jets, which are seen at a small angle to the line of sight, similarly to blazars. This claim was further strengthened by the *Fermi* satellite discovery of high-energy emission from a handful of RLNLS1 sources (e.g. Foschini 2011; Karamanavis 2015; Yao et al. 2015, and references therein). Further support of their blazar-like nature came from the Monitoring Of Jets in Active galactic nuclei with VLBA Experiments (MOJAVE, Lister et al. 2009) survey. Among the five *Fermi*-detected RLNLS1 sources included in their sample, three show jet components moving at superluminal speeds (Lister et al. 2016; see also Fuhrmann et al. 2016).

Thus, flat-spectrum RLNLS1 sources are thought to be similar to blazars, but residing (mostly) in spiral host galaxies with central black holes of lower masses (Crenshaw, Kraemer & Gabel 2003). A statistical study by Berton et al. (2015) showed that the black hole mass distribution is the same for steep-spectrum and flat-spectrum RLNLS1 sources. The suggested scenario of Berton et al. (2015) is that at larger inclination angle, instead of flat-spectrum RLNLS1 sources, one starts to see steep-spectrum RLNLS1 sources. At even larger inclination angles, the Doppler effect broadens the line in the disc-like shaped broad-line regions (e.g. Shen & Ho 2014), and a broad-line radio galaxy hosted by a disc galaxy can be observed. Finally, when looking through the obscuring torus, the objects are described as narrow-line radio galaxies hosted by disc galaxies.

The multiwavelength study of Järvelä, Lähteenmäki & León-Tavares (2015) showed that the jet is the main source of the radio, optical and X-ray emission in RLNLS1 sources, while in radio-quiet NLS1 sources, the infrared and radio emission mostly originate from star formation. The origin of infrared emission in RLNLS1 sources however is unclear, it is possible that star formation and reradiated emission from the torus also contribute (Caccianiga et al. 2015; Järvelä et al. 2015).

SDSS J110006.07+442144.3 (hereafter, J1100+4421) is a newly discovered NLS1 candidate source. It was found in the Kiso Supernova Survey (Morokuma et al. 2014) by Tanaka et al. (2014), thanks to its dramatic optical flare on 2014 February 23.

Follow-up observations of Tanaka et al. (2014) revealed that the FWHMs of the broad components of Mg II and H β lines in J1100+4421 are smaller than 2000 km s^{-1} . The redshift of the source was measured to be $z = 0.84$. It was classified as an NLS1 candidate. Tanaka et al. (2014) calculated the black hole mass, $\sim 1.5 \times 10^7 M_{\odot}$, using the Mg II line, and $\sim 1.0 \times 10^7 M_{\odot}$, using the H β line. They also calculated the bolometric luminosity ($6 \times 10^{44} \text{ erg s}^{-1}$), which is 30 per cent of the Eddington luminosity. These values are consistent with the findings of Foschini et al. (2015), who analysed the multiwavelength properties of 42 RLNLS1s, and found Eddington ratios of 1 per cent to 49 per cent, and black hole masses in the range of 10^6 – $10^8 M_{\odot}$. However, J1100+4421 did not show a Fe II bump, and the flux ratio of the [O III] line to the H β is much larger (~ 5 – 9) than the value used to define NLS1 sources. According to Tanaka et al. (2014), the former can be attributed to the flaring state of the source, which can cause the emergence of strong continuum emission. With respect to the latter, the luminosity of the [O III] line was shown to correlate with the radio power (Labiano 2008). Thus, Tanaka et al. (2014) argue that the especially luminous line in J1100+4421 may be related to strong jet emission in the source. The presence of jet is also indicated by the extreme radio-loudness of J1100+4421. The ratio of the 1.4-GHz flux density measured in the Faint Images of the Radio Sky at Twenty-Centimeters (FIRST) survey (Becker, White & Helfand 1995) to the quiescent flux density measured at $4400 \text{ \AA}$ is

~ 3000 . When the optical flux density is measured at the highest point of the flaring state, this ratio is still ~ 400 (Tanaka et al. 2014).

To image the radio structure of the source at milliarcsecond (mas) resolution and ascertain whether it has a blazar-like jet emission, we observed J1100+4421 with very long baseline interferometry (VLBI) using the European VLBI Network (EVN) at 1.7 and 5 GHz. We also searched for a possible γ -ray counterpart to J1100+4421 in the latest *Fermi*/Large Array Telescope (LAT) data. In the following, we assume a flat Λ cold dark matter cosmological model with $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.27$ and $\Omega_{\Lambda} = 0.73$. At the redshift of the source ($z = 0.84$), 1-arcsec angular size corresponds to 7.754 kpc projected linear size (Wright 2006).

2 OBSERVATIONS AND DATA REDUCTION

2.1 EVN data

The EVN observations of J1100+4421 took place on 2015 February 10 at 1.7 GHz and on 2015 March 24 at 5 GHz. At 1.7 GHz, the interferometric array consisted of seven antennas: Effelsberg (Germany), the Jodrell Bank Mark 2 telescope (the United Kingdom), Medicina (Italy), Onsala (Sweden), Toruń (Poland), the Westerbork Synthesis Radio Telescope (WSRT, the Netherlands) and Sheshan (China). At 5 GHz, Noto (Italy) and Yebes (Spain) were also added to the array. However, Medicina did not observe and the WSRT produced no useful data at the higher frequency.

The observations were carried out in e-VLBI mode (Szomoru 2008). The signals received at the radio telescopes were transmitted over optical fibre networks directly to the central data processor for real-time correlation. The correlation with 2 s integration time was done at the EVN software correlator (SEFXC, Keimpema et al. 2015), in the Joint Institute for VLBI ERIC (JIVE), Dwingeloo, the Netherlands. Eight intermediate-frequency channels (IFs) were used in both polarizations at both bands. Each IF had a width of 16 MHz and was divided into 32 spectral channels. The total bandwidth was 256 MHz. The observations were performed in phase-reference mode (Beasley & Conway 1995). The target and the phase calibrator were observed alternately, with ~ 3.5 min spent on the target and ~ 1 min spent on the calibrator. The phase-reference calibrator used at both frequencies was J1108+4330, separated by $\sim 1^\circ 7'$ from the target in the sky. Its coordinates are right ascension $\alpha_{\text{cal}} = 11^{\text{h}}08^{\text{m}}23^{\text{s}}.47694$ and declination $\delta_{\text{cal}} = +43^\circ 30' 53''.6571$, the uncertainties are 0.2 mas in both directions.¹ Additionally, 4C39.25 was included as fringe-finder at both bands. The observations lasted for 4.5 and 4.25 h, and on-source integration times were 164 and 119 min, at 1.7 and 5 GHz, respectively.

The data were reduced in the standard manner (Diamond 1995) using the U.S. National Radio Astronomy Observatory (NRAO) Astronomical Image Processing System (AIPS, Greisen 2003). First, interferometric visibility amplitudes were calibrated using the gain curves and system temperature measurements obtained at the telescope sites. Then, fringe-fitting was performed on the phase-calibrator and fringe-finder sources. Their visibility data were exported to be imaged with the DIFMAP program package (Shepherd, Pearson & Taylor 1994). The hybrid mapping procedure was used with several cycles of CLEANING (Högbom 1979) and phase self-calibration. Gain correction factors were determined in DIFMAP and subsequently were used in AIPS to scale the amplitudes. The average

¹ Data are obtained from <http://astrogeo.org> maintained by L. Petrov, rfc2015b solutions.

Figure 1. Naturally weighted 1.7-GHz EVN image of J1100+4421. Observation took place on 2015 February 10. The peak brightness is $19.4 \text{ mJy beam}^{-1}$, the restoring beam is $24.8 \text{ mas} \times 8.2 \text{ mas}$ (FWHM) at a position angle of 9° and shown in the lower left-hand corner of the image. The lowest contours are at $\pm 3.5\sigma$ noise level ($\pm 0.1 \text{ mJy beam}^{-1}$), further positive contour levels increase by a factor of 2. The dashed line represents negative contour. The labels denote the Gaussian components fitted to the visibility data.

corrections factors did not exceed 10 per cent. Four–four channels, the most affected by the bandpass response function, were discarded at the beginning and end of each IF. Then fringe-fitting on the calibrator was repeated in AIPS, now taking into account its brightness distribution using its CLEAN component model obtained in DIFMAP. The derived solutions were interpolated and applied to the target source, J1100+4421. The hybrid mapping of the target was also performed in DIFMAP. No amplitude self-calibration was attempted on the source, and phase self-calibration was not done for Sheshan for time intervals shorter than 60 min at 1.7 GHz and 15 min at 5 GHz.

Since J1100+4421 was sufficiently bright and compact, we also performed fringe-fitting directly on the target source at both frequencies. Then, we imaged the source in the same way as described above. The images obtained (Figs 1 and 2) were in good agreement with those obtained by transferring the phase solutions of the calibrator. Comparing the peak brightness values in the two images in both bands, the coherence loss in phase-referencing was 1 per cent at 1.7 GHz and 14 per cent at 5 GHz.

2.2 Archival VLA data

J1100+4421 was also observed during a 40-s snapshot at 8.4 GHz with the Very Large Array (VLA) in its most extended A configuration on 1995 August 15 (project code: AM484) within the Cosmic Lens All-Sky Survey (Myers et al. 2003). The data were downloaded from the NRAO archive² and analysed following standard data reduction steps in AIPS. The absolute flux density calibrator

Figure 2. Naturally weighted 5-GHz EVN image of J1100+4421. Observation took place on 2015 March 24. The peak brightness is $23.2 \text{ mJy beam}^{-1}$, the restoring beam is $3.8 \text{ mas} \times 1.2 \text{ mas}$ (FWHM) at a position angle of 16° and shown in the lower left-hand corner of the image. The lowest contours are at $\pm 3.5\sigma$ noise level ($\pm 0.2 \text{ mJy beam}^{-1}$), further positive contour levels increase by a factor of 2. The dashed line represents negative contour. The label denotes the Gaussian component fitted to the visibility data.

Figure 3. 8.4-GHz VLA image of J1100+4421. The observation was performed on 1995 August 15. The peak brightness is $18.4 \text{ mJy beam}^{-1}$. The FWHM size of the restoring beam is $0.27 \times 0.25 \text{ arcsec}^2$ at a position angle of -81° , as shown in the bottom left-hand corner of the image. The lowest contours are at $\pm 0.5 \text{ mJy beam}^{-1}$ corresponding to 3σ image noise level, further positive contour levels increase by a factor of 2.

used in the project was 3C286. The calibrated visibility data were exported from AIPS, imaging was done in DIFMAP. The resulting map is shown in Fig. 3. The weights of the data points are set inversely proportional to the amplitude errors (natural weighting).

² <http://archive.nrao.edu>

Table 1. Details of the EVN observations of J1100+4421 (right ascension $\alpha_t = 11^{\text{h}}00^{\text{m}}06^{\text{s}}.0571$, declination $\delta_t = +44^{\circ}21'44''.383$). In Columns 2, 3, 4, 5 and 6, the parameters of the fitted circular Gaussian components are listed. In Columns 7, 8, 9 and 10, the peak intensity, the size and position angle of the restoring beam and the rms noise levels of the radio images shown in Figs 1 and 2 are given.

ν (GHz)	ID	Relative position		Flux density (mJy)	FWHM (mas)	peak mJy beam ⁻¹	beam size (mas × mas)	PA (°)	rms (mJy beam ⁻¹)
		RA (mas)	Dec. (mas)						
1.7	C	–	–	20.9 ± 0.1	2.5 ± 0.2	19.4	24.8×8.2	9	0.029
1.7	N	-12.5 ± 2.0	79 ± 2.0	1.5 ± 0.3	16.0 ± 4.0	–	–	–	–
5	C	–	–	25.0 ± 0.4	0.38 ± 0.05	23.2	3.8×1.2	16	0.057

2.3 Fermi LAT data

J1100+4421 has no associated counterpart in the newest and so far most comprehensive γ -ray catalogue the *Fermi*/LAT 4-Year Point Source Catalog (3FGL, Acero et al. 2015). The closest 3FGL source is 3FGL J1105.7+4427 with a separation of more than 60 arcmin.

We analysed data from LAT using the *Fermi*/LAT Science Tools v10r0p5³ software package. The LAT data were extracted in a circle with 15° radius around the coordinates of J1100+4421, over the whole energy range, choosing only LAT data type ‘photon’. We used the unbinned likelihood analysis and followed the provided tutorial⁴ in the data reduction. The parameters used for different tasks were the ones suggested in the tutorial. First, only events of class ‘source’ in the energy band of 100 MeV–100 GeV were chosen using `gtselect`, and a zenith angle cut of 90° was implemented to remove the disturbance caused by the Earth’s limb. Next `gtmktime` was used to select only the good time intervals for further analysis. An exposure map was generated using `gtlcube` and `gtexpmap`. The newest galactic diffuse emission model (`gll_iem_v06.fits`) and the extragalactic isotropic diffuse emission model (`iso_P8R2_SOURCE_V6_v06.txt`), which includes the residual cosmic-ray background, were used for computing the diffuse source responses with `gtdiffrrps`. Finally, J1100+4421 and all the sources within 10° radius (29 sources) were modelled with `gtlike` using the `powerlaw2` model and the unbinned likelihood algorithm with the optimizer MINUIT.

We first integrated over the whole time period from 2008 August 5 00:00:00 UTC to 2016 December 19 00:00:00 UTC (3058 d) for which we have data. There are some very faint sources nearby, having test statistic (TS) values less than 0, causing the fit not to converge. We kept removing the sources with $TS < 0$, except J1100+0044, and running the reduction again, until the TS values of all nearby sources were positive. Over the whole period, the TS value for J1100+0044 was negative, meaning that the fit without the source is better than with it, implying that it is very improbable that this source has emitted detectable γ -rays during this period. To verify this result, we integrated over the same time period in bins of 180 d, resulting in 17 bins, using the same parameters and procedure. The TS value for J1100+4421 was negative in every bin, further indicating that J1100+4421 has not been active at γ -rays between 2008 August and 2016 December. The upper limit of the γ -ray flux is 9.9×10^{-10} photon cm⁻² s⁻¹ in the 100 MeV–300 GeV energy band.

3 RESULTS

3.1 The pc-scale radio structure

A single, compact radio-emitting feature was detected at 5 GHz in our EVN observations (Fig. 2). At 1.7 GHz, an additional feature is detected north from the central source (Fig. 1). The phase-referencing observations allowed us to obtain the accurate coordinates of J1100+4421. We used the `MAXFIT` verb in `AIPS` to determine the right ascension and declination of the brightness peak at the 5-GHz EVN image: $\alpha_t = 11^{\text{h}}00^{\text{m}}06^{\text{s}}.0571$, $\delta_t = +44^{\circ}21'44''.383$. The coordinates are the same within the uncertainties at the two observing frequencies. We estimate that the obtained coordinates are accurate within 1 mas.

We used `DIFMAP` to fit the visibilities with brightness distribution models. At 1.7 GHz, two circular Gaussian components are needed (labelled C and N), while at 5 GHz, a single circular Gaussian component (C) adequately modelled the visibilities. The parameters of the fitted components (projected separations from the core, flux densities and FWHM sizes) are summarized in Table 1.

Using the size and flux density values derived from the higher frequency measurement, one can calculate the brightness temperature as

$$T_B = 1.22 \times 10^{12} (1+z) \frac{S}{\theta^2 \nu^2} \text{ K}, \quad (1)$$

where z is the redshift of the source, S is the flux density given in Jy, θ is the FWHM size in mas and ν is the observing frequency in GHz. The obtained brightness temperature is $(1.56 \pm 0.4) \times 10^{10}$ K. However, if the FWHM size derived from the model fit is smaller than the smallest resolvable size of the array, the component is unresolved, and the brightness temperature value is only a lower limit. According to Kovalev et al. (2005), the minimum resolvable FWHM size of a Gaussian component in our observation at 5 GHz would be 0.18 mas along the major axis of the restoring beam, indicating that the feature is not unresolved. However, the more recent work of Natarajan et al. (2017) showed that calibration uncertainties have significant effect on the minimum resolvable size. Investigating synthetic observations made with an array very similar to our EVN observation, they concluded that the minimum resolvable size can be a factor of 2 larger, ~ 0.4 mas. Therefore, the fitted FWHM size of the component is an upper limit, and the derived brightness temperature of J1100+4421 can be regarded as a lower limit only.

The recovered flux density of the core component was somewhat larger at 5 GHz than at 1.7 GHz. Formally, the compact source has a spectral index of $\alpha = 0.20 \pm 0.03$ ($S \sim \nu^\alpha$). However, we note that the EVN observations were not simultaneous but separated by six weeks. Compact radio-emitting AGNs are known to show variations in the radio on time-scales of weeks to

³ <http://fermi.gsfc.nasa.gov/ssc/data/analysis/software/>

⁴ http://fermi.gsfc.nasa.gov/ssc/data/analysis/scitools/likelihood_tutorial.html

Figure 5. Non-contemporaneous radio flux density measurements of J1100+4421. Black circles show data from various surveys such as the Giant Meterwave Radio Telescope 150 MHz all-sky radio survey (TGSS, Intema et al. 2017), the Westerbork Northern Sky Survey (WENSS, Rengelink et al. 1997), the NVSS, the FIRST and the Green Bank 4.85 GHz survey (GB6, Gregory et al. 1996), and the flux density from the archival 8.4-GHz VLA observation (project code: AM484). Red squares (grey squares in the printed version) show the sum of the flux density of the fitted components of our EVN data.

would expect to see a morphology reminiscent of a more resolved source rather than a single compact feature.

Tanaka et al. (2014) pointed out that the observed optical variability is similar to that of the γ -ray detected RLNLS1 sources. They compared the spectral energy distribution of J1100+4421 to the γ -ray-loud RLNLS1s of PMN J0948+0022, and PKS 2004–447 (Abdo et al. 2009a,b), and hypothesized that assuming the same photon index as that of PMN J0948+0022, the upper limit of its γ -ray luminosity would be comparable to the measured γ -ray luminosity of PMN J0948+0022. However, our analysis of all available *Fermi*/LAT data resulted in no detection of γ -rays at the position of J1100+4421.

Morokuma et al. (2017) conducted extensive monitoring of J1100+4421 in the optical and near-infrared bands. They showed that based upon the spectral shape, the optical and near-infrared emission of the source is dominated by emission from the jet. From the fastest observed variations, and assuming a Doppler factor of 10, they estimate that the region responsible for the variability must be smaller than 5.4×10^{15} cm.

4.2 The large-scale radio structure

In Fig. 5, the radio flux density measurements of J1100+4421 are summarized. The data are not simultaneous, and the observations are conducted with different angular resolutions, therefore source variability as well as resolution issues may complicate the picture. Nevertheless, up until 1.4 GHz, the spectrum seems to follow a steep power law, which might flatten afterwards. The red symbols represent the total flux density recovered in our high resolution EVN observations. The 1.7-GHz data point clearly shows the source variability (as discussed in Section 3.2), while the discrepancy between the two 5-GHz data points can also be caused by the different resolutions of the observations [the EVN observation recovered only the pc-scale core emission, while in the Green Bank 4.85-GHz radio survey (Gregory et al. 1996), the kpc-scale structure could have been measured as well], as well as source variability.

The kpc-scale radio emission around J1100+4421 can only be seen in the FIRST image. In the 8-GHz VLA image, there was no additional radio emission above $0.8 \text{ mJy beam}^{-1}$, 4.5σ image noise level in a $30 \times 30 \text{ arcsec}^2$ region around the source. This may indicate that the extended emission has a steep spectrum between 1.4 and 8.4 GHz. The circular Gaussian component describing the northern feature detected at 1.7 GHz in our EVN observation is at $\sim 80 \text{ mas}$, corresponding to $\sim 620 \text{ pc}$ projected distance from the core. This feature may be the jet connecting to the northern lobe seen in the FIRST image. There is no sign of any other radio emission down to a noise level of $0.13 \text{ mJy beam}^{-1}$ (4σ) at a region of $8 \times 8 \text{ arcsec}^2$ around the centre in the EVN image. If the two lobes contained compact features, the sensitivity of our EVN observation would have been adequate to detect them. Therefore, they are not compact, their sizes have to be larger than the largest recoverable size of the interferometer (Wrobel 1995), which is $\sim 70 \text{ mas}$ in the 1.7-GHz EVN observation, corresponding to $\sim 540 \text{ pc}$ at the redshift of J1100+4421.

The morphology of the large-scale structure of J1100+4421 can be classified as two-sided. Doi et al. (2012) calculated the core dominance parameter for six RLNLS1 with kpc-scale radio structures as the flux density ratio between the core and lobes measured in the FIRST images. To disentangle the emission from the lobe and core region, we fitted the central region of the FIRST image of J1100+4421 with a circular Gaussian component. To keep this feature compact, we used a fixed size of $5.4 \times 5.4 \text{ arcsec}^2$, the clean beam size of the image. The resulting flux density of this unresolved component is $11.0 \pm 0.2 \text{ mJy}$. Then, we used the AIPS verb IMSTAT to measure the flux density of the lobes in the residual image. They each have flux densities of $\sim 3 \text{ mJy}$. Thus, the core dominance parameter is 1.8 for J1100+4421, the smallest among the core-dominated sources in Doi et al. (2012). Following Doi et al. (2012), we can estimate the jet kinetic energy in the large-scale radio structure of J1100+4421. The 1.4-GHz radio power of a $\sim 3 \text{ mJy}$ flux density lobe in J1100+4421 assuming a spectral index of -0.7 is $\sim 9 \times 10^{24} \text{ W Hz}^{-1}$. According to the scaling relation of Cavagnolo et al. (2010), the jet kinetic power is thus $3 \times 10^{37} \text{ W}$. This value is slightly higher than the minimum value, $\sim 10^{37} \text{ W}$ estimated to be necessary to create supersonic lobes according to Doi et al. (2012). Thus, the radio power in the extended structure of J1100+4421 is intermediate between the radio powers in FR I and FR II radio galaxies (Fanaroff & Riley 1974).

Currently, there are a handful of RLNLS1 sources reported to have extended, kpc-scale radio structure (Whalen et al. 2006; Antón, Browne & Marchã 2008; Gliozzi et al. 2010; Doi et al. 2012; Richards & Lister 2015; Congiu et al. 2017), the projected size ranging between ~ 23 and $\sim 100 \text{ kpc}$. Additionally, recently Yao et al. (2015) reclassified a flat-spectrum quasar, SDSS J122222.55+041315.7 as an NLS1 source. According to its FIRST image and the VLA measurement at 1.4 GHz reported by Kharb, Lister & Cooper (2010), this RLNLS1 also exhibits a significant extended radio emission, with a projected linear size of $\sim 160 \text{ kpc}$. Thus, the extended radio emission seen in J1100+4421 (Fig. 4), with a projected size of $\sim 150 \text{ kpc}$ at 1.4 GHz, is among the largest of the known RLNLS1 sources.

Assuming similar kpc-scale velocities as the ones for the six RLNLS1 sources investigated by Doi et al. (2012), it would require $\sim 10^6 \text{ yr}$ to build the radio morphology seen in the FIRST image. However, this is a lower limit only, since the viewing angle of the structure is not known, therefore its deprojected size might be larger. On the other hand, the similar flux density values seen in both lobes

may indicate that the structure is oriented close to the plane of the sky.

Morokuma et al. (2017) report on the detection of a faint galaxy at a distance of 2.7 to the east from J1100+4421. We did not detect any radio counterpart at the position of this galaxy down to 0.5 and 0.1 mJy beam⁻¹, 3 σ level from the VLA 8-GHz data and our EVN 1.7-GHz data, respectively. However, it seems unlikely, it cannot be completely ruled out that the large-scale radio structure detected in the FIRST observation is partly related to this galaxy.

5 SUMMARY

The source J1100+4421 was discovered during a dramatic optical flare by Tanaka et al. (2014). Based upon the follow-up optical observations, which showed unusually narrow broad lines, the source was classified as a candidate NLS1 source. However, the two other defining characteristics of NLS1 sources (presence of Fe II bump and low flux of [O III] λ 5007 compared to H β line) do not apply to J1100+4421. Nevertheless, the derived low black hole mass and relatively high Eddington rate are in agreement with the proposed possible NLS1 nature of the source (Cracco et al. 2016). Tanaka et al. (2014) suggested that the optical flare is most probably caused by the jet emission, and the optical and near-infrared monitoring campaign of Morokuma et al. (2017) showed that both the optical and near-infrared emissions are dominated by synchrotron radiation from the jet.

Our high-resolution EVN observations showed that the source indeed has a mas-scale compact radio structure, similar to other very radio-loud NLS1 sources. The derived lower limit on the brightness temperature is 1.56×10^{10} K, which is below but still consistent with the equipartition limit, therefore we cannot exclude the presence of Doppler boosting in the source. At lower resolution, J1100+4421 shows an extended, two-sided radio morphology of ~ 150 kpc, which is among the largest of the known RLNLS1 sources. Somewhat contrary to the results of the VLBI observations, the large-scale radio emission indicates that the radio structure is oriented close to the plane of the sky. So far the source has not been detected in the *Fermi*/LAT γ -ray mission.

To reveal the connection and possible jet bending between the inner, compact radio-emitting feature and the large-scale lobes, high-sensitivity, intermediate-resolution radio interferometric observations are desirable. Comparing the flux densities measured in different radio observations, it is clear that J1100+4421 shows significant variability at cm wavelengths at least on time-scales of years. If the source shows radio variability on shorter time-scales, frequent radio flux density monitoring could provide independent estimate of the Doppler factor.

ACKNOWLEDGEMENTS

We thank the referee for careful reading of the manuscript and for his/her comments, which helped to improve the paper. KEG was supported by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences. We wish to thank Venkatesh Ramakrishnan for the help with the *Fermi* data reduction. The European VLBI Network is a joint facility of independent European, African, Asian and North American radio astronomy institutes. Scientific results from data presented in this publication are derived from the following EVN project code: EG087. The e-VLBI research infrastructure in Europe was supported by the European Community's Seventh Framework Programme (FP7/2007-2013) under grant agreement

RI-261525 NEXPreS. The research leading to these results has received funding from the European Commission Seventh Framework Programme (FP/2007-2013) under grant agreement no. 283393 (RadioNet3). This research was supported by the Hungarian National Research Development and Innovation Office (OTKA NN110333) and the China–Hungary Collaboration and Exchange Programme by the International Cooperation Bureau of the Chinese Academy of Sciences (CAS). TA acknowledges the grant of the Youth Innovation Promotion Association of CAS.

REFERENCES

- Abdo A. A. et al., 2009a, *ApJ*, 699, 976
- Abdo A. A. et al., 2009b, *ApJ*, 707, L142
- Ajero F. et al., 2015, *ApJS*, 218, 23
- Angelakis E. et al., 2015, *A&A*, 575, A55
- Antón S., Browne I. W. A., Marchã M. J., 2008, *A&A*, 490, 583
- Beasley A. J., Conway J. E., 1995, in Zensus J. A., Diamond P. J., Napier P. J., eds, *ASP Conf. Ser. Vol. 82, Very Long Baseline Interferometry and the VLBA*. Astron. Soc. Pac., San Francisco, p. 327
- Becker R. H., White R. L., Helfand D. J., 1995, *ApJ*, 450, 559
- Berton M. et al., 2015, *A&A*, 578, A28
- Caccianiga A. et al., 2015, *MNRAS*, 451, 1795
- Cavagnolo K. W., McNamara B. R., Nulsen P. E. J., Carilli C. L., Jones C., Birzan L., 2010, *ApJ*, 720, 1066
- Collin S., Kawaguchi T., 2004, *A&A*, 426, 797
- Condon J. J., Cotton W. D., Greisen E. W., Yin Q. F., Perley R. A., Taylor G. B., Broderick J. J., 1998, *AJ*, 115, 1693
- Congiu E. et al., 2017, *A&A*, 603, A32
- Cracco V., Ciroi S., Berton M., Di Mille F., Foschini L., La Mura G., Rafanelli P., 2016, *MNRAS*, 462, 1256
- Crenshaw D. M., Kraemer S. B., Gabel J. R., 2003, *AJ*, 126, 1690
- Diamond P. J., 1995, in Zensus J. A., Diamond P. J., Napier P. J., eds, *ASP Conf. Ser. Vol. 82, Very Long Baseline Interferometry and the VLBA*. Astron. Soc. Pac., San Francisco, p. 227
- Doi A., Nagira H., Kawakatu N., Kino M., Nagai H., Asada K., 2012, *ApJ*, 760, 41
- Fanaroff B. L., Riley J. M., 1974, *MNRAS*, 167, 31P
- Foschini L., 2011, in Foschini L., Colpi M., Gallo L., Grupe D., Komossa S., Leighly K., Mathur S., eds, *Proceedings of Science, Vol. NLS1, Narrow-Line Seyfert 1 Galaxies and their Place in the Universe*. POS, Trieste, p. 24
- Foschini L. et al., 2015, *A&A*, 575, A13
- Fuhrmann L. et al., 2016, *Res. Astron. Astrophys.*, 16, 176
- Giozzi M., Papadakis I. E., Grupe D., Brinkmann W. P., Raeth C., Kedziora-Chudczer L., 2010, *ApJ*, 717, 1243
- Gregory P. C., Scott W. K., Douglas K., Condon J. J., 1996, *ApJS*, 103, 427
- Greisen E. W., 2003, in Heck A., ed, *Information Handling in Astronomy – Historical Vistas*. Kluwer, Dordrecht, p. 109
- Gu M., Chen Y., Komossa S., Yuan W., Shen Z., Wajima K., Zhou H., Zensus J. A., 2015, *ApJS*, 221, 3
- Helfand D. J., White R. L., Becker R. H., 2015, *ApJ*, 801, 26
- Högbom J. A., 1979, *A&AS*, 36, 173
- Homan D. C. et al., 2006, *ApJ*, 642, L115
- Intema H. T., Jagannathan P., Mooley K. P., Frail D. A., 2017, *A&A*, 598, A78
- Järvelä E., Lähteenmäki A., León-Tavares J., 2015, *A&A*, 573, A76
- Karamanavis V., 2015, PhD thesis, Max-Planck-Institut für Radioastronomie
- Keimpema A. et al., 2015, *Exp. Astron.*, 39, 259
- Kellermann K. I., Sramek R., Schmidt M., Shaffer D. B., Green R., 1989, *AJ*, 98, 1195
- Kharb P., Lister M. L., Cooper N. J., 2010, *ApJ*, 710, 764
- Komossa S., Voges W., Xu D., Mathur S., Adorf H.-M., Lemson G., Duschl W. J., Grupe D., 2006, *AJ*, 132, 531
- Kovalev Y. Y. et al., 2005, *AJ*, 130, 2473
- Labiano A., 2008, *A&A*, 488, L59

- Lister M. L. et al., 2009, *AJ*, 138, 1874
 Lister M. L. et al., 2016, *AJ*, 152, 12
 Mathur S., 2000, *MNRAS*, 314, L17
 Morokuma T. et al., 2014, *PASJ*, 66, 114
 Morokuma T. et al., 2017, *Publ. Astron. Soc. Japan*, 69, 82
 Myers S. T. et al., 2003, *MNRAS*, 341, 1
 Natarajan I., Paragi Z., Zwart J., Perkins S., Smirnov O., van der Heyden K., 2017, *MNRAS*, 464, 4306
 Osterbrock D. E., Pogge R. W., 1985, *ApJ*, 297, 166
 Pogge R. W., 2000, *New Astron. Rev.*, 44, 381
 Readhead A. C. S., 1994, *ApJ*, 426, 51
 Rengelink R. B., Tang Y., de Bruyn A. G., Miley G. K., Bremer M. N., Roettgering H. J. A., Bremer M. A. R., 1997, *A&AS*, 124
 Richards J. L., Lister M. L., 2015, *ApJ*, 800, L8
 Shen Y., Ho L. C., 2014, *Nature*, 513, 210
 Shepherd M. C., Pearson T. J., Taylor G. B., 1994, *BAAS*, 26, 987
 Szomoru A., 2008, *Proc. Sci., EXPRes and the e-EVN. SISSA, Trieste, PoS(IX EVN Symposium)040*
 Tanaka M. et al., 2014, *ApJ*, 793, L26
 Whalen D. J., Laurent-Muehleisen S. A., Moran E. C., Becker R. H., 2006, *AJ*, 131, 1948
 Wright E. L., 2006, *PASP*, 118, 1711
 Wrobel J. M., 1995, in Zensus J. A., Diamond P. J., Napier P. J., eds, *ASP Conf. Ser. Vol. 82, Very Long Baseline Interferometry and the VLBA. Astron. Soc. Pac., San Francisco*, p. 411
 Yao S., Yuan W., Zhou H., Komossa S., Zhang J., Qiao E., Liu B., 2015, *MNRAS*, 454, L16
 Yuan W., Zhou H. Y., Komossa S., Dong X. B., Wang T. G., Lu H. L., Bai J. M., 2008, *ApJ*, 685, 801
 Zhou H., Wang T., Yuan W., Lu H., Dong X., Wang J., Lu Y., 2006, *ApJS*, 166, 128
 Zhou H. et al., 2007, *ApJ*, 658, L13

This paper has been typeset from a $\text{\LaTeX}$ file prepared by the author.