This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Mai-Gisondi, Galina; Maaheimo, Hannu; Chong, Sun-Li; Hinz, Sandra; Tenkanen, Maija; Master, Emma

Functional comparison of versatile carbohydrate esterases from families CE1, CE6 and CE16 on acetyl-4-O-methylglucuronoxylan and acetyl-galactoglucomannan

Published in:

BIOCHIMICA ET BIOPHYSICA ACTA: GENERAL SUBJECTS

DOI:

[10.1016/j.bbagen.2017.06.002](https://doi.org/10.1016/j.bbagen.2017.06.002)

Published: 01/09/2017

Document Version

Peer reviewed version

Published under the following license:

CC BY-NC-ND

Please cite the original version:

Mai-Gisondi, G., Maaheimo, H., Chong, S-L., Hinz, S., Tenkanen, M., & Master, E. (2017). Functional comparison of versatile carbohydrate esterases from families CE1, CE6 and CE16 on acetyl-4-O-methylglucuronoxylan and acetyl-galactoglucomannan. *BIOCHIMICA ET BIOPHYSICA ACTA: GENERAL SUBJECTS*, 1861(9), 2398-2405. <https://doi.org/10.1016/j.bbagen.2017.06.002>

Functional comparison of versatile carbohydrate esterases from families CE1, CE6 and CE16 on acetyl-4-*O*-methylglucuronoxylan and acetyl-galactoglucomannan

Galina Mai-Gisondi¹, Hannu Maaheimo², Sun-Li Chong³, Sandra Hinz⁴, Maija Tenkanen⁵,
Emma Master^{1,6*}

¹Department of Chemical Technology and Biotechnology, Aalto University; FI-00076 Aalto, Kemistintie 1, Espoo, Finland

²VTT Technical Research Centre of Finland Ltd, P.O. Box 1000, FI-02044 VTT Espoo, Finland

³Department of Food and Environmental Sciences, University of Helsinki; FI-00014 University of Helsinki, Latokartanonkaari 11, Helsinki, Finland
moved to: Department of Biology and Biological Engineering Division of Industrial Biotechnology, Chalmers University of Technology, Kemivägen 10, SE-412 96 Göteborg, Sweden

⁴DuPont Industrial Biosciences, Nieuwe Kanaal 7-S, 6709 PA Wageningen, The Netherlands

⁵Department of Food and Environmental Sciences, University of Helsinki; FI-00014 University of Helsinki, Latokartanonkaari 11, Helsinki, Finland

⁶Department of Chemical Engineering and Applied Chemistry, University of Toronto, 200 College Street, Toronto, Ontario, M5S 3E5, Canada

*Corresponding author:

Emma R. Master

Telephone: 416-946-7891

Fax: 416-978-8605

Email: emma.master@utoronto.ca

Keywords: Acetyl (xylan) esterases, carbohydrate esterase families, hemicellulose, acetyl-4-*O*-methylglucuronoxylan and acetyl-galactoglucomannan, regio-selectivity

ABSTRACT

Background: The backbone structure of many hemicelluloses is acetylated, which presents a challenge when the objective is to convert corresponding polysaccharides to fermentable sugars or else recover hemicelluloses for biomaterial applications. Carbohydrate esterases (CE) can be harnessed to overcome these challenges.

Methods: Enzymes from different CE families, *AnAcXE* (CE1), *OsAcXE* (CE6), and *MtAcE* (CE16) were compared based on action and position preference towards acetyl-4-*O*-methylglucuronoxylan (MGX) and acetyl-galactoglucomannan (GGM). To determine corresponding positional preferences, the relative rate of acetyl group released by each enzyme was analyzed by real time ^1H NMR.

Results: *AnAcXE* (CE1) showed lowest specific activity towards MGX, where *OsAcXE* (CE6) and *MtAcE* were approximately four times more active than *AnAcXE* (CE1). *MtAcE* (CE16) was further distinguished by demonstrating 100 times higher activity on GGM compared to *AnAcXE* (CE1) and *OsAcXE* (CE6), and five times higher activity on GGM than MGX. Following 24 h incubation, all enzymes removed between 78 to 93% of total acetyl content from MGX and GGM, where *MtAcE* performed best on both substrates.

Major Conclusions: Considering action on MGX, all esterases showed preference for doubly substituted xylopyranosyl residues (2,3-*O*-acetyl-Xylp). Considering action on GGM, *OsAcXE* (CE6) preferentially targeted 2-*O*-acetyl-mannopyranosyl residues (2-*O*-acetyl-Manp) whereas *AnAcXE* (CE1) demonstrated highest activity towards 3-*O*-acetyl-Manp positions; regiopreference of *MtAcE* (CE16) on GGM was less clear.

General significance: The current comparative analysis identifies options to control the position of acetyl group release at initial stages of reaction, and enzyme combinations likely to accelerate deacetylation of major hemicellulose sources.

1. INTRODUCTION

Hemicelluloses constitute approximately 20–30% of the total mass of annual and perennial plants. They are differentiated based on backbone and side group chemistries, which vary depending on the botanical source [1,2]. 4-*O*-methylglucuronoxylans (MGX) are the main hemicellulose in deciduous wood such as eucalyptus trees, and contain xylopyranosyl (Xylp) backbone sugars that are connected through β -(1→4)-linkages and are substituted by α -(1→2)-linked 4-*O*-methylglucopyranosyluronic acid (MeGlc_pA) (Figure 1). By contrast, galactoglucomannans (GGM) are the main hemicellulose in coniferous wood such as spruce trees, and contain β -(1→4)-linked D-glucopyranosyl (Glc_p) and β -D-mannopyranosyl (Man_p) units that are likewise connected through β -(1→4) glycosidic bonds (Figure 1). The galactoglucomannan backbone is also substituted by α -(1→6)-linked-D-galactose side groups, mainly to Man_p in spruce galactoglucomannans [3].

Glucuronoxylans and galactoglucomannans are both acetylated in their natural forms [2]. Several biological functions have been attributed to the acetylation of these main hemicelluloses, including protection of plant cell walls from degradation by microbial enzymes, along with regulated interaction with cellulose [4,5]. Acetylation may occupy single O-2 or O-3 positions of Xylp or Man_p subunits in glucuronoxylans and galactoglucomannans, respectively. Xylp in glucuronoxylans may also be di-acetylated [6–8], and Xylp decorated with 2-*O*-MeGlc_pA can be acetylated at the O-3 position ((2-*O*-MeGlc_pA)3-*O*-acetyl-Xylp) [6,7,9]. A summary of reported compositions of acetyl-4-*O*-methylglucuronoxylans and acetyl-galactoglucomannans is provided in Supplemental Table 1.

The carbohydrate active enzyme (CAZy) classification system (www.cazy.org) [10] currently assigns carbohydrate esterases (CEs) into 16 CE families. Deacetylating enzymes with reported activity on xylans and/or corresponding oligosaccharides include acetyl xylan esterases (AcXE; EC. 3.1.1.72) belonging to CE families CE1-7, an unclassified carbohydrate esterase [11], as well as acetyl esterases (AcE; EC. 3.1.1.6) belonging to family CE16 [12–14]. Nevertheless, only AXEs from CE1, CE3-CE6, and CE16 were confirmed to include enzymes with preferred activity towards xylans [15, 42]. By contrast, comparatively few enzymes have been reported to deacetylate galactoglucomannans or corresponding oligosaccharides. Notable examples include: 1) an AcE from *Trichoderma reesei* VTT-D-86271 (Rut C 30) (TrCE16) [16–19] an acetyl glucomannan esterase (AGME) and feruloyl esterase (FE; CE1) from *Aspergillus oryzae* VTT-D-85248 (presently renamed as *Aspergillus tubingensis*) [18–20]; 3) an esterase within a commercial

enzyme preparation from *Aspergillus niger* (Celluzyme) [21,22] and 4) two AcXEs belonging to CE1, one from *Penicillium purpurgenum* [23] and other from *Schizophyllum commune* VTT-D-88362 [24]. Low levels of acetyl group release from oligomers of galactoglucomannans was also detected in culture filtrates of *Aspergillus awamori* VTT-D-71025, *Aureobasidium pullulans* VTT-D-89397 and *Streptomyces olivochromogens* VTT-E-82157 [18].

Considering the impacts of hemicellulose acetylation summarized above, AcEs and AcXEs can be harnessed to promote hemicellulose saccharification [25–27], control the rheology and solubility of hemicelluloses [1,13], and improve thermomechanical pulp (TMP) yield [28]. The applied significance of AcEs and AcXEs, along with the inability to predict enzyme action based on CAZy family assignation alone, has motivated functional comparisons of several CE families [8,13,29–32]. Neumüller et al. [32] propose three distinguishing groups of xylan-active CEs: 1) those acting only on O-2 and O-3 monoacetylated Xylp, which have been identified in families CE2, CE3, and CE4; 2) those with further activity towards 2,3-di-O-acetylated Xylp, which have been identified in families CE1, CE5 and CE6, and 3) those with further activity towards 3-O acetylated Xylp substituted at O-2 by MeGlcAp; which have been identified in family CE16 (e.g. *An*CE16). Puchart et al. [33] subsequently isolated aldo-tetraouronic acid ($\text{Ac}^3\text{MeGlcA}^3\text{Xyl}_3$) to compare the ability of selected CE16s to target 3-O acetylated Xylp substituted also at O-2 by MeGlcAp. Notably, *Tr*CE16 was the only CE16 esterase able to release acetyl group from both isomers of $\text{Ac}^3\text{MeGlcA}^3\text{Xyl}_3$, containing acetyl group at O-3 and O-4, where acetyl group migration from O-3 to O-4 is possible.

Whereas recent comparative analyses of CEs have considerably advanced our understanding of enzyme regio-selectivity towards acetylated xylans, only one study has been reported that investigates CE action towards specific acetylated positions within mannan substrates [24]. Accordingly, herein we compare the activity and regio-selectivity of three CEs using major, wood-derived hemicelluloses, namely 4-O-methylglucuronoxylan (MGX) (herein, from *Eucalyptus globulus*) and acetyl-galactoglucomannan (GGM) (herein, from Norway spruce). The CEs were selected from three different CE families, namely acetyl xylan esterase *An*AcXE (CE1) from *Aspergillus nidulans*, acetyl xylan esterase *Os*AcXE (CE6) from *Orpinomyces* sp., and acetyl esterase *Mt*AcE (CE16) from *Myceliophthora thermophila*. Enzymes from these CE families were selected based on (1) our previous analysis of *An*AcXE (CE1) that confirmed activity on acetylated polysaccharides, including cellulose acetate [34], (2) *Os*AcXE (CE6) being a commercially available enzyme with reported activity on acetylated xylans [29,32], but so far not on mannans, which is notable given that the CE6 family comprises the largest number of

predicted plant acetyl esterases and so may be expected to possess broad substrate specificity and act on both xylans and (gluco)mannans, and (3) few examples where CE16 enzymes have been characterized using natural substrates even though reports to date indicate activity towards typically resistant positions, such as (2-*O*-MeGlc₄pA)3-*O*-acetyl-Xyl₆p in MGX [33], and reported activity towards (oligomeric) GGM [14,17,18].

Briefly, the current comparative analysis showed that positional preference of selected enzymes towards MGX does not predict its positional preference towards GGM, and that enzymes showing similar positional preference towards one substrate may differ when compared using another. The present comparative analysis of CEs on both MGX and GGM underscores the importance of including natural substrates when characterizing enzyme action, and the difficulty to predict enzyme action on GGM based on known activity towards MGX or vice versa.

2. MATERIALS AND METHODS

2.1. Enzymes, substrates, and assay reagents

In-house and commercially available enzymes were used for this study. Specifically, *AnAcXE* (AN6093.2) from *Aspergillus nidulans* was recombinantly produced in *Pichia pastoris* at pH 5.0 and 30 °C and then purified based on [35]. The CE6 *AcXE* from *Orpinomyces* sp. (AAC14690.1; *OsAcXE*) was purchased from Megazyme (Wicklow, Ireland). The CE16 *AcE* acetyl esterase from *Myceliophthora thermophila* (AGW01024.1; *MtAcE*) [8] was produced at DuPont Biosciences (Wageningen, Netherlands) and polished using size exclusion and hydrophobic chromatography (Supplemental Figure 1).

Acetyl-4-*O*-methylglucuronoxylan (MGX) isolated from milled (8 mm) chips of eucalyptus by steam extraction [36] was kindly provided by Prof. J.C. Parajó (University of Vigo, Spain) and acetyl-galactoglucomannan (GGM) obtained from the TMP mill process waters [37] was from Prof. Willför (Åbo Akademi University, Finland). Total acetyl groups available in MGX or GGM were determined as described in [9] and were found to be 15 and 9% of total substrate weight, respectively. Briefly, 1 mg of GGM or MGX was suspended in 200 µL of 0.1 N NaOH and incubated with shaking (120 rpm) for 24 h at room temperature; released acetic acid was neutralized and then measured using the Acetic Acid Assay Kit (K-ACET) purchased from Megazyme (Ireland).

2.2. pH Optima

1 The pH optimum of each enzyme was evaluated using 4-methylumbelliferyl acetate (4-MUA)[38]
2 (Supplemental Figure 2). Reaction mixtures (400 μ L) comprised 100 mM of the selected buffer,
3 2.5 mM 4-MUA, and 10 μ L of the enzyme sample. Enzyme doses were adjusted to ensure that
4 linear rates of reaction were measured, and were 3.6 μ g for *AnAcXE* (CE1); 0.2 μ g for *OsAcXE*
5 (CE6) and 0.05 μ g for *MtAcE* (CE16). Chosen buffers were: sodium citrate buffer (pH 3.0 to pH
6 5.0), sodium phosphate buffer (pH 6.0 to pH 8.0), and glycine-OH buffer (pH 9.0 to pH 10.0).
7 Following incubation at 40 °C for 10 min, reactions were stopped by adding of 600 μ L 50 mM
8 citric acid (pH 2.2), vortexed, and then passed through a 0.2 μ m GHP Acrodiscs 13 filter (PALL)
9 to remove insoluble particles prior to measurement of 4-methylumbelliferone at 345 nm. pH
10 optima were also confirmed using MGX and the Acetic Acid kit (K-ACET) (Supplemental Figure
11 3).

12

13 **2.3. Enzyme stability at pH 6.0**

14

15 Because overnight reactions with MGX and GGM were performed at pH 6.0 (see below),
16 enzyme stabilities were also evaluated at this pH condition. In this case, 2.5 μ g of *AnAcXE*
17 (CE1), *OsAcXE* (CE6), or *MtAcE* (CE16) were incubated for 24 h in 50 μ L of 100 mM sodium
18 phosphate buffer (pH 6.0). Following the incubation period, 3.6 μ g of *AnAcXE* (CE1), 0.2 μ g of
19 *OsAcXE* (CE6), and 0.05 μ g of *MtAcE* (CE16) were transferred to 50 μ L reaction mixtures
20 comprising 100 mM sodium phosphate buffer (pH 6.0) and 2.5 mM 4-MUA. Reaction products
21 were then processed and measured as described above.

22

23 **2.4. Activity measurements using MGX and GGM**

24

25 To measure specific activities for each enzyme on each substrate, reaction mixtures (200 μ L)
26 comprised 100 mM sodium phosphate buffer (pH 7.0), 0.5% (w/v) substrate, and 50 μ L of the
27 enzyme sample. Enzyme doses were adjusted to ensure that linear rates of reaction were
28 measured. When using MGX, enzyme doses were 0.5, 1 or 5 μ g of *AnAcXE* (CE1), 0.5, 1 or 5
29 μ g of *OsAcXE* (CE6), and 0.5 or 1 μ g of *MtAcE* (CE16). When using GGM, enzyme doses were
30 2.5 or 5 μ g of *AnAcXE* (CE1), 2.5 or 5 μ g of *OsAcXE* (CE6), and 0.05, 0.1 or 0.3 μ g of *MtAcE*.
31 Following incubation at 40 °C for 10 min, reactions were stopped by adding 40 μ L of 0.33 M
32 H₂SO₄. Released acetic acid was quantified using the Acetic Acid Assay Kit (K-ACET).

33

34 To quantify the extent of acetyl group released by each enzyme after 24 h, reaction mixtures (200
35 μ L) comprising 100 mM sodium phosphate buffer (pH 6.0), 0.5% (w/v) substrate, and 10 μ g of

AnAcXE (CE1), *OsAcXE* (CE6), or *MtAcE* (CE16) (i.e., 10 mg enzyme/ g substrate) were incubated with shaking (120 rpm) at 40 °C. Overnight reactions were performed at pH 6.0 to minimize the possibility of auto hydrolysis, release of acetyl groups from the substrate, and acetyl group migration [39]. Reaction mixtures without enzyme served as negative controls and were subtracted from test measurements.

2.5. HSQC NMR spectroscopy

Quantitative HSQC NMR was performed to quantify acetyl group release from MGX and GGM. Analyses of MGX were carried out in D₂O, whereas analyses of GGM were carried out in DMSO-d₆. All NMR spectra were collected on a 600 MHz Bruker Avance III NMR spectrometer equipped with a QCI H-C/N/P-D cryogenically cooled probe head. The measurements were carried out at either 22 °C (samples dissolved in D₂O) or 60 °C (samples dissolved in DMSO-d₆). For the 1D ¹H spectra the residual water signal was suppressed by a four second volume selective presaturation (so called 1D-NOESY presaturation). (Neuhaus et al. 1996). The quantitative HSQC spectra were acquired using matched sweep adiabatic pulses optimised for ¹³C sweep width of 130 – 10 ppm for all 180° ¹³C pulses in order to compensate the differences in the ¹J_{CH} coupling constants (Bruker's pulse program hsqcetgpsi2.3) (Zwahlen et al. 1997). Matrices of 2048 x 256 data points were collected and zero filled once in F1; a $\pi/2$ shifted squared sine bell weighting function was applied in both dimensions prior to the Fourier transformation.

2.6. HSQC NMR Spectra annotations and quantifications

The chemical shifts for MGX were referenced to the C-1 and H-1 signal of MeGlc₆P₆A (5.28 ppm, 98.85 ppm); chemical shifts for GGM in D₂O were referenced to C-1 of Glc₆P₆A (103.55 ppm) and H-2 of 2-*O*-acetyl Man₆P₆A (5.42 ppm). In the case of GGM in DMSO, the chemical shifts were calibrated against DMSO-d₆ (2.50 ppm, 39.51 ppm). Annotation of MGX spectra were according to [7] and [6]; annotation of GGM spectra were according to [40].

The relative content of the acetylated Xyl₆P in the MGX was calculated from quantitative heteronuclear single quantum coherence (qHSQC) spectra according to [41]. Briefly, the signals of the H-1 and C-1 of substituted and nonsubstituted Xyl₆P were summed to 100%, thereafter the signals of H-2 of 2-*O*-acetyl, H-3 of 3-*O*-acetyl, H-2 of 2,3-*O*-acetyl, and H-3 of (2-*O*-MeGlc₆P₆A)3-*O*-acetyl-Xyl₆P were integrated separately to calculate the relative content of each

form of *O*-acetyl-Xylp subunit. For the quantitation of relative content of acetylated Manp in the GGM, the H-1 and C-1 of the substituted and non-substituted Manp, as well as the Glcp and Galp were summed as 100%, thereafter the signal of H-2 of 2-*O*-acetyl- and H-3 of 3-*O*-acetyl-Manp was integrated separately to calculate the relative content of each form of *O*-acetyl Manp to 100%, thereafter the signals of H-2 of 2-*O*-acetyl, H-3 of 3-*O*-acetyl, H-2 of 2,3-*O*-acetyl, and H-3 of (2-*O*-MeGlc(A)3-*O*-acetyl-Xylp were integrated separately to calculate the relative content of each form of *O*-acetyl-Xylp subunit. For the quantitation of relative content of acetylated Manp in the GGM, the H-1 and C-1 of the substituted and non-substituted Manp, as well as the Glcp and Galp were summed as 100%, thereafter the signal of H-2 of 2-*O*-acetyl- and H-3 of 3-*O*-acetyl-Manp was integrated separately to calculate the relative content of each form of *O*-acetyl Manp.

2.7. ¹H NMR Analyses

Rates of acetyl group release from specific positions within MGX and GGM were monitored by ¹H NMR. Both substrates (600 mg) were dissolved in 600 μL of 100 mM sodium phosphate buffer (pH 6.0) following replacement of milliQ water by D₂O through freeze drying. All samples were analyzed in 5.0 mm NMR tubes (Aldrich) using a Bruker Avance III 400 MHz spectrometer equipped with a 5 mm BBFO Plus probe head. A water suppression pulse program (noesygppr1d) with suppression power of 3.1623e-006 W was used for relative quantitative measurements. The following acquisition parameters were applied: 90° pulse with relaxation delay of 4 s and acquisition time of 5.1 s; 8 scans and 4 dummy scans; 65536 data points and spectrum width of 15.979 ppm. Before the reaction was initiated by addition of enzyme, the spectra of native substrates were measured at 40 °C. After enzyme addition, acquisition parameters were adjusted (e.g., to align signals) and then spectra were automatically recorded every 1.8 min over 30 min (Figure 2). When using MGX, enzyme doses per mg of substrate were 5 μg of *AnAcXE* (CE1), 1 μg of *OsAcXE* (CE6), and 1 μg of *MtAcE* (CE16). When using GGM, enzyme doses per mg of substrate were 10 μg of *AnAcXE* (CE1), 10 μg *OsAcXE* (CE6), and 1 μg/mg *MtAcE* (CE16). In all cases, enzymes were prepared in 100 mM deuterated sodium phosphate buffer, spectra were processed and analyzed using TopSpin 3.0 (Bruker).

Sum of integrals was normalized to 100% and signals were then plotted against time using Origin 2016 64 bit (Supplemental Figure 4), and resulting slopes were used to calculate the rate of acetyl groups release from specific positions by each enzyme [32].

3. RESULTS and DISCUSSION

3.1. Establishing reaction conditions for comparative analyses

AnAcXE (CE1), *OsAcXE* (CE6), and *MtAcE* (CE16) were similarly active at pH 7.0 or pH 6.0 (Table 1), by showing $\geq 90\%$ of their maximal activity at pH 7.0 and $\geq 85\%$ at pH 6.0. Accordingly, reaction rates were measured at pH 7.0 to ensure comparable substrate solubility, whereas overnight reactions were performed at pH 6.0 to minimize the possibility of autohydrolysis, non-enzymatic release of acetyl groups, and acetyl group migration. Notably, *AnAcXE* and *MtAcE* retained over 90% activity after 24 h at pH 6.0 (40 °C), whereas under these conditions, *OsAcXE* activity decreased by 40% after 5 h and 60% after 24 h (Figure 3).

3.2. Extent of carbohydrate esterase action on MGX

Acetic acid measurement and HSQC NMR analyses were performed to compare the deacetylation efficiency of *AnAcXE* (CE1), *OsAcXE* (CE6), and *MtAcE* (CE16) on MGX. In general, slightly higher deacetylation efficacies were calculated from HSQC spectra as compared to acetic acid measurements (Figure 4; Table 2), which likely reflects the relative sensitivity of the analytical methods used.

Following 24 h of incubation, maximal removal of acetyl groups from MGX by *AnAcXE* (CE1) was between 80%-90% (Figure 4; Table 2). This result was similar to that reported for other CE1 acetyl xylan esterases, including: (1) a CE1 from *Schizophyllum commune* (~80% deacetylation of DMSO-extracted birchwood MGX) [24]; (2) *AnAXE* from *Aspergillus niger* and *MtAXE3* from *Myceliophthora thermophila* C1 (76-88% deacetylation of *O*-acetylated neutral xylo-oligosaccharides and 50–60% deacetylation of *O*-acetylated MeGlcA-substituted xylo-oligosaccharides [8]); and (3) *TeCE1* from *Talaromyces emersonii* (~80% deacetylation of MeGlcA-substituted xylooligosaccharides) [32].

Similar to *AnAcXE* (CE1), *OsAcXE* (CE6) released 80-90% of acetyl groups from MGX (Figure 4; Table 2). Earlier analyses of *OsAcXE* (also known as *OsCE6*) show less than 50% of total acetyl group release from MGX. However, in that case, MGX enriched in acidic xylo-oligosaccharides AcUXOS (i.e., (2-*O*-MeGlcA)-3-*O*-acetyl-Xylp) was used [32], and

Koutaniemi et al. [8] previously showed that enzymatic deacetylation of acidic xylo-oligosaccharides is typically lower than neutral xylo-oligosaccharides. Compared to *AnAcXE* (CE1) and *OsAcXE* (CE6), *MtAcE* (CE16) released the highest extent of acetyl groups from all positions in MGX (~90% total released). By comparison, *AnCE16* from *Aspergillus niger* was reported to release 18% of acetyl groups from AcUXOS derived from *Eucalyptus globulus* [32], whereas *TrCE16* from *Trichoderma reesei* was reported to release 10% of acetyl groups from xylo-oligomers with DP of ~10 [16,17]. These results suggested that *TrCE16* and *AnCE16* activity is restricted to the non-reducing end of corresponding substrates [19,32]. Different to both *AnCE16* and *TrCE16*, *PaCE16* was able to act on birchwood acetyl-glucuronoxylan leading to substrate precipitation [42]. Thus, given the extent of acetyl groups released from MGX measured in the current study, *MtAcE* (CE16) likely targets both non-reducing and internal positions within targeted substrates and so in this regard, is more similar to *PaCE16* than *TrCE16* or *AnCE16*.

Quantitative HSQC (qHSQC) was then used to identify acetylated positions in MGX most susceptible to enzyme action. Less than 1% of 2-*O*-acetyl-Xylp positions, and approximately 5% of 3-*O*-acetyl-Xylp and 2,3-*O*-acetyl-Xylp positions remained following MGX treatment with *AnAcXE* (CE1). Similarly, less than 5% 2-*O*-acetyl-Xylp positions in MGX remained intact following treatment of MGX with *OsAcXE* (CE6). Somewhat lower deacetylation efficiencies were measured for *OsAcXE* (CE6) action towards 3-*O*-acetyl-Xylp and 2,3-*O*-acetyl-Xylp positions, where 10% and 20% of corresponding acetyl groups remained.

By comparison, *MtAcE* (CE16) effectively targeted the broadest range of acetyl group positions, leaving between 3-5% of 2-*O*-acetyl-Xylp, 3-*O*-acetyl-Xylp, and 2,3-*O*-acetyl-Xylp positions, while also partially removing acetyl groups from (2-*O*-MeGlcA)-3-*O*-acetyl-Xylp positions (Table 2; Supplemental Figure 4B). Low but detectable and reproducible activity of *MtAcE* towards (2-*O*-MeGlcA)-3-*O*-acetyl-Xylp after 24 h of reaction is consistent with reports of other CE16 enzymes, including *P. anserina* (*PaCE16A*) [42], which shares approximately 70% of protein sequence identity with *MtAcE* (CE16), but also *TrCE16* from *T. reesei* [14,33,43] and *AnCE16* from *A. niger* [32,33]. It was earlier postulated [8] and later shown [33] that acetyl group migration from O-3 to O-4 could account, at least partially, for *MtAcE* (CE16) activity towards (2-*O*-MeGlcA)-3-*O*-acetyl-Xylp.

3.3. Extent of carbohydrate esterase action on GGM

1 All three enzymes were able to act on GGM in addition to MGX. During the 24 h incubation,
2 *MtAcE* released approximately 90% of acetyl groups in GGM, whereas roughly 80% of acetyl
3 groups were released by *AnAcXE* (CE1) and *OsAcXE* (CE6) under the same conditions
4 (Figure 4). *MtAcE* (CE16) action on GGM was comparable to that of an acetyl mannan esterase
5 (AGME) from *Aspergillus niger* [20,21], an acetyl xylan esterase (CE1) from *Schizophyllum*
6 *commune* [24], and a feruloyl esterase (CE1) from *Aspergillus oryzae* [24]. In each of these
7 cases, between 90 and 95% of acetyl groups from GGM were released. By comparison, the
8 family CE1 AcXEs from *Penicillium purpurogenum* (30%) and AGME (unclassified CE) from
9 *Aspergillus oryzae* [24] were previously shown to release approximately 70% of acetyl groups
10 from GGM.

11 Consistent with the expected higher degree of polymerization of GGM (recovered from a TMP
12 process waters) compared to MGX (recovered from a steam explosion process), reaction products
13 from GGM precipitated upon enzyme treatment. As a result, qHSQC analyses could not be
14 performed in D₂O or DMSO-d₆. Accordingly, ¹H NMR spectra were collected to identify acetyl
15 group positions most resistant to enzyme action (Figure 5). Most notably, *OsAcXE* (CE6) targeted
16 all acetyl groups in GGM, whereas 2(b)-*O*-acetyl-Manp was resistant to hydrolysis by *AnAcXE*
17 (CE1) and *MtAcE* (CE16). Positions 2(a)-*O*-acetyl-Manp and 2(b)-*O*-acetyl-Manp (along with 3(a)-
18 *O*-acetyl-Manp and 3(b)-*O*-acetyl-Manp) differ in terms of neighboring sugars [40,44]. Retention of
19 2(b)-*O*-acetyl-Manp in GGM therefore reveals that neighboring groups along the polysaccharide
20 backbone influence enzyme accessibility to pendent acetyl groups.

21

22 **3.4. Specific activity and positional preference of selected carbohydrate esterases**

23

24 *OsAcXE* (CE6) and *MtAcE* (CE16) showed similar specific activity towards MGX, which were
25 approximately four times higher than that obtained for *AnAcXE* (CE1) (Table 3). By contrast,
26 similar specific activities were measured for *AnAcXE* (CE1) and *OsAcXE* (CE6) on GGM, which
27 were lower than corresponding enzyme activities on MGX (Table 3). *MtAcE* (CE16) was
28 interestingly distinguished by demonstrating 100 times higher activity on GGM compared to
29 *AnAcXE* (CE1) and *OsAcXE* (CE6), and five times higher activity on GGM than MGX (Table 3).
30 The specific activity of *MtAcE* (CE16) towards GGM (~2000 nkat/mg; 120 µg/min/mg) was
31 comparable to that reported for AGME from *Aspergillus niger* (1190 nkat/mg) where 5 times
32 higher GGM concentration was used [20,21]. Similarities between *MtAcE* (CE16) and AGME
33 suggests that *MtAcE* (CE16) is likewise an unspecific acetyl mannan esterase [8].

34

35 Real time ¹H NMR was then performed for each enzyme to unravel potential preferences for

specific positions within MGX and GGM. Positional preference was defined as the ratio of deacetylation rate for a given acetyl group position to the abundance of corresponding acetyl groups in the original substrate. It is important to note here that calculated rates of acetyl group release from 2-*O*-acetyl-Xylp and 3-*O*-acetyl-Xylp may underestimate true values since reaction products after removal of one of the acetyl groups from 2,3-*O*-acetyl-Xylp will contribute to these signals.

Although the specific activity of *AnAcXE* on MGX differed from both other enzymes, *AnAcXE* (CE1), *OsAcXE* (CE6) and *MtAcE* (CE16) similarly demonstrated comparatively high activity towards acetyl groups present on doubly substituted Xylp (i.e., 2,3-*O*-acetyl-Xylp, Table 4).

Although it was not possible to resolve which of these two acetyl residues were preferentially targeted, preferred action towards 2,3-*O*-acetyl-Xylp positions was previously also reported for *Talaromyces emersonii* *TeCE1* [32], *Orpinomyces* *sp. PC-2* *OsCE6* [29,32], and *Aspergillus niger* *AnCE16A* [33].

Considering singly substituted Xylp positions, *AnAcXE* (CE1), *OsAcXE* (CE6) and *MtAcE* (CE16) all revealed preference towards 2-*O*-acetyl-Xylp positions, which is consistent with reported activities for above mentioned *TeCE1*, *OsCE6* and *AnCE16A* [29,32,33]. *AnAcXE* (CE1) and *OsAcXE* (CE6) were similarly active towards 3-*O*-acetyl-Xylp, whereas activity towards this position was not observed for *MtAcE* (CE16) (Table 4). In contrast to *MtAcE* (CE16), *TrCE16* was shown to preferentially target 3-*O*-acetyl-Xylp and 4-*O*-acetyl-Xylp positions of oligosaccharides of acetyl-glucuronoxylan [43]. However, *AnCE16A* rapidly targets 2,3-*O*-acetyl-Xylp positions, followed by 2-*O*-acetyl-Xylp and 3-*O*-acetyl-Xylp positions, of polymeric xylans [33], but poorly targets 2-*O*-acetyl-Xylp positions in methyl β -D-xylopyranoside diacetates and triacetates (Puchart et al. 2016), thus showing differing position specificity with polymeric and monomeric substrates. Similarly, then, the difference in reported positional preference of *MtAcE* and *TrCE16* may be due to differences in the length of substrates used in corresponding analyses.

Whereas *AnAcXE* (CE1) and *OsAcXE* (CE6) displayed similar positional preferences in MGX, the positional preferences of *OsAcXE* (CE6) and *MtAcE* (CE16) were most similar when evaluated using GGM. Specifically, when considering the anomeric region of corresponding ¹H NMR spectra where peak signals were clearly resolved, it could be seen that both *OsAcXE* (CE6) and *MtAcE* (CE16) displayed similar preference for 2-*O*-acetyl-Manp positions in GGM, whereas highest rates of *AnAcXE* (CE1) activity were towards 3-*O*-acetyl-Manp substituents. Notably, however, positional preference within GGM was less clear when considering the acetyl region of corresponding ¹H NMR spectra, where peak signals are more intense but also overlapping (Figure

2B, Table 5). Still, the comparably high activity of *MtAcE* (CE16), as well as *AnAcXE* (CE1) and *OsAcXE* (CE6) towards the O-2 acetylated position of both *Xylp* in MGX and *Manp* in GGM suggests that stereochemistry plays a minor role in substrate recognition by these enzymes. *TrCE16* preferentially targets 3-*O*-acetyl-*Xylp* in oligo-saccharides of MGX; it was also shown to deacetylate oligomers of GGM [18], however, regio-selective activity of *TrCE16* or other AcEs from CE16 on oligomers or polymers of GGM has not been characterized.

4. CONCLUSION

The comparison of three CE families using both MGX and GGM uncovered substrate-dependent and enzyme dependent differences in reaction rates, extent of substrate conversion, and regio-selectivity. In particular, the acetyl xylan esterases *AnAcXE* (CE1) and *OsAcXE* (CE6) displayed different specific activities towards MGX yet similar regio-selectivity. On the other hand, these enzymes were similarly active towards GGM. Notably, *MtAcE* (CE16) was set apart from both acetyl xylan esterases by its comparatively high specific activity towards GGM. Nevertheless, comparably high activity of all three enzymes on 2-*O*-acetylated positions in GMX and GGM, which has equatorial orientation in *Xylp* and axial orientation in *Manp*, suggests that the stereochemistry of the acetyl group has little effect on the activity of these enzymes.

The comparative analysis of three CE families on MGX and GGM underscore the impact of the selected substrate on reported enzyme activity as well as regio-selectivity, further highlighting known challenges associated with predicting enzyme action based on model compounds. Positions within major hemicellulose sources that remain resistant to CE action were confirmed, including (2-*O*-MeGlc_pA)3-*O*-acetyl-*Xylp* in MGX, and some 2-*O*-acetyl-*Manp* or 3-*O*-acetyl-*Manp* in GGM, thereby identifying targets for enzyme discovery as well as enzyme combinations that could be harnessed to promote hemicellulose recovery (e.g., through precipitation) versus full saccharification. Finally, earlier reports which tested AXE activity on ground wood powder [45], and fungal AXE expressed in plants [26], show that enzyme action on hemicelluloses embedded with plant fibre could be predicted from enzyme action on extracted hemicelluloses. However, direct comparison of different AXEs on MGX or GGM present in plant fibre remains to be done.

ACKNOWLEDGEMENTS

This work was supported by a grant to GM from Ella and Georg Ehrnrooth foundation (Finland), a FiDiPro Fellowship to ERM from the Finnish Funding Agency for Technology and Innovation (Tekes), and an ERC Consolidator Grant to EM (BHIVE – 648925).

REFERENCES

- [1] A. Ebringerová, Z. Hromádková, T. Heinze, Hemicellulose, *Adv. Polym. Sci.* 186 (2005) 1–67. doi:10.1007/b136816.
- [2] H. V Scheller, P. Ulvskov, Hemicelluloses, *Annu. Rev. Plant Biol.* 61 (2010) 263–289. doi:10.1146/annurev-arplant-042809-112315.
- [3] S. Willför, R. Sjöholm, C. Laine, M. Roslund, J. Hemming, B. Holmbom, Characterisation of water-soluble galactoglucomannans from Norway spruce wood and thermomechanical pulp, *Carbohydr. Polym.* 52 (2003) 175–187. doi:10.1016/S0144-8617(02)00288-6.
- [4] P.M.-A. Pawar, S. Koutaniemi, M. Tenkanen, E.J. Mellerowicz, Acetylation of woody lignocellulose: significance and regulation, *Front. Plant Sci.* 4 (2013) 118. doi:10.3389/fpls.2013.00118.
- [5] M. Busse-Wicher, T.C.F. Gomes, T. Tryfona, N. Nikolovski, K. Stott, N.J. Grantham, D.N. Bolam, M.S. Skaf, P. Dupree, The pattern of xylan acetylation suggests xylan may interact with cellulose microfibrils as a twofold helical screw in the secondary plant cell wall of *Arabidopsis thaliana*, *Plant J.* 79 (2014) 492–506. doi:10.1111/tpj.12575.
- [6] D. V. Evtuguin, J.L. Tomás, A.M.S. Silva, C.P. Neto, Characterization of an acetylated heteroxylan from *Eucalyptus globulus* Labill, *Carbohydr. Res.* 338 (2003) 597–604. doi:10.1016/S0008-6215(02)00529-3.
- [7] A. Teleman, J. Lundqvist, F. Tjerneld, H. Stålbrand, O. Dahlman, Characterization of acetylated 4-O-methylglucuronoxylan isolated from aspen employing ¹H and ¹³C NMR spectroscopy, *Carbohydr. Res.* 329 (2000) 807–15. doi:10.1016/S0008-6215(00)00249-4.
- [8] S. Koutaniemi, M.P. van Gool, M. Juvonen, J. Jokela, S.W. Hinz, H.A. Schols, M. Tenkanen, Distinct roles of carbohydrate esterase family CE16 acetyl esterases and polymer-acting acetyl xylan esterases in xylan deacetylation, *J. Biotechnol.* 168 (2013) 684–92. doi:10.1016/j.jbiotec.2013.10.009.
- [9] A. Teleman, M. Tenkanen, A. Jacobs, O. Dahlman, Characterization of O-acetyl-(4-O-methylglucurono)xylan isolated from birch and beech, *Carbohydr. Res.* 337 (2002) 373–377. doi:10.1016/S0008-6215(01)00327-5.
- [10] V. Lombard, H.G. Ramulu, E. Drula, P.M. Coutinho, B. Henrissat, The carbohydrate-active enzymes database (CAZy) in 2013, *Nucleic Acids Res.* 42 (2014) D490–D495.

doi:10.1093/nar/gkt1178.

- [11] O. Alalouf, Y. Balazs, M. Volkinshtein, Y. Grimpel, G. Shoham, Y. Shoham, A new family of carbohydrate esterases is represented by a GDSL hydrolase/acetylxylan esterase from *Geobacillus stearothermophilus*, *J. Biol. Chem.* 286 (2011) 41993–42001. doi:10.1074/jbc.M111.301051.
- [12] P. Biely, J. Puls, H. Schneider, Acetyl xylan esterases in fungal cellulolytic systems, *FEBS Lett.* 186 (1985) 80–84. doi:10.1016/0014-5793(85)81343-0.
- [13] P. Biely, Microbial carbohydrate esterases deacetylating plant polysaccharides, *Biotechnol. Adv.* 30 (2012) 1575–1588. doi:10.1016/j.biotechadv.2012.04.010.
- [14] F.A. Adesioye, T.P. Makhalanyane, P. Biely, D.A. Cowan, Phylogeny, classification and metagenomic bioprospecting of microbial acetyl xylan esterases, *Enzyme Microb. Technol.* 93–94 (2016) 79–91. doi:10.1016/j.enzmictec.2016.07.001.
- [15] P. Biely, B. Westereng, V. Puchart, P. De Maayer, D.A. Cowan, Recent Progress in Understanding the Mode of Action of Acetylxylan Esterases, 44 (2014). doi:10.5458/jag.jag.
- [16] K. Poutanen, M. Sundberg, An acetyl esterase of *Trichoderma reesei* and its role in the hydrolysis of acetyl xylans, *Appl. Microbiol. Biotechnol.* 28 (1988) 419–424. doi:10.1007/BF00268207.
- [17] K. Poutanen, M. Sundberg, H. Korte, J. Puls, Deacetylation of xylans by acetyl esterases of *Trichoderma reesei*, *Appl. Microbiol. Biotechnol.* 33 (1990) 506–510. <http://www.springerlink.com/index/m171047h67876585.pdf> (accessed January 18, 2013).
- [18] M. Tenkanen, J. Puls, M. Rättö, L. Viikari, Enzymatic deacetylation of galactoglucomannans, *Appl. Microbiol. Biotechnol.* 39 (1993) 159–165. doi:10.1007/BF00228600.
- [19] M. Tenkanen, Action of *Trichoderma reesei* and *Aspergillus oryzae* esterases in the deacetylation of hemicelluloses, *Biotechnol. Appl. Biochem.* 27 (1998) 19–24. doi:10.1111/j.1470-8744.1998.tb01370.x.
- [20] M. Tenkanen, J. Thornton, L. Viikari, An acetylglucomannan esterase of *Aspergillus oryzae*; purification, characterization and role in the hydrolysis of O-acetyl-galactoglucomannan, *J. Biotechnol.* 42 (1995) 197–206. doi:10.1016/0168-1656(95)00080-A.
- [21] J. Puls, Acetylmannanesterase: a new component in the arsenal of wood mannan degrading enzymes, in: M. Kuwahara, M. Shimada (Eds.), *Biotechnol. Pulp Pap. Ind.*, Uni Publishers Co.Ltd, Tokyo, 1992: pp. 357–363.
- [22] R.P. de Vries, J. Visser, *Aspergillus* enzymes involved in degradation of plant cell wall polysaccharides, *Microbiol. Mol. Biol. Rev.* 65 (2001) 497–522. doi:10.1128/MMBR.65.4.497.

- 1 [23] R. Gutiérrez, E. Cederlund, L. Hjelmqvist, a Peirano, F. Herrera, D. Ghosh, W. Duax, H.
2 Jörnvall, J. Eyzaguirre, Acetyl xylan esterase II from *Penicillium purpurogenum* is similar to
3 an esterase from *Trichoderma reesei* but lacks a cellulose binding domain., *FEBS Lett.* 423
4 (1998) 35–8. <http://www.ncbi.nlm.nih.gov/pubmed/9506837>.
- 5 [24] M. Tenkanaan, J. Eyzaguirre, R. Isoniemi, C.B. Faulds, P. Biely, Comparison of catalytic
6 properties of acetyl xylan esterases from three carbohydrate esterase families, in: *Am. Chem.*
7 *Soc.*, 2003: pp. 211–229.
- 8 [25] H. Rabemanantsoa, S. Saka, Various pretreatments of lignocellulosics, *Bioresour. Technol.*
9 199 (2016) 83–91. doi:10.1016/j.biortech.2015.08.029.
- 10 [26] P.M.A. Pawar, M. Derba-Maceluch, S.L. Chong, L.D. Gómez, E. Miedes, A. Banasiak, C.
11 Ratke, C. Gaertner, G. Mouille, S.J. Mcqueen-Mason, A. Molina, A. Sellstedt, M. Tenkanen,
12 E.J. Mellerowicz, Expression of fungal acetyl xylan esterase in *Arabidopsis thaliana*
13 improves saccharification of stem lignocellulose, *Plant Biotechnol. J.* 14 (2016) 387–397.
14 doi:10.1111/pbi.12393.
- 15 [27] J. Zhang, M. Siika-Aho, M. Tenkanen, L. Viikari, The role of acetyl xylan esterase in the
16 solubilization of xylan and enzymatic hydrolysis of wheat straw and giant reed, *Biotechnol.*
17 *Biofuels.* 4 (2011) 1–9. doi:10.1186/1754-6834-4-60.
- 18 [28] J. Thornton, M. Tenkanen, R. Ekman, B. Holmbom, L. Viikari, Possibility of increasing
19 mechanical pulp yield by enzymatic treatment, *Holzforschung.* 48 (1994) 436–440.
- 20 [29] I. Uhliaríková, M. Vršanská, B. V McCleary, P. Biely, Positional specificity of acetyl xylan
21 esterases on natural polysaccharide: an NMR study, *Biochim. Biophys. Acta.* 1830 (2013)
22 3365–3372. doi:10.1016/j.bbagen.2013.01.011.
- 23 [30] L. Pouvreau, M.C. Jonathan, M.A. Kabel, S.W.A. Hinz, H. Gruppen, H.A. Schols,
24 Characterization and mode of action of two acetyl xylan esterases from *Chrysosporium*
25 *lucknowense* C1 active towards acetylated xylans, *Enzyme Microb. Technol.* 49 (2011) 312–
26 320. doi:10.1016/j.enzmictec.2011.05.010.
- 27 [31] D.L. Blum, X.-L. Li, H. Chen, L.G. Ljungdahl, Characterization of an acetyl xylan esterase
28 from the anaerobic fungus *Orpinomyces* sp. strain PC-2, *Appl. Environ. Microbiol.* 65 (1999)
29 3990–3995.
30 [http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=99731&tool=pmcentrez&rendert](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=99731&tool=pmcentrez&rendertype=abstract)
31 [ype=abstract](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=99731&tool=pmcentrez&rendertype=abstract).
- 32 [32] K.G. Neumüller, A.C. de Souza, J.H. van Rijn, H. Streekstra, H. Gruppen, H.A. Schols,
33 Positional preferences of acetyl esterases from different CE families towards acetylated 4-O-
34 methyl glucuronic acid-substituted xylo-oligosaccharides, *Biotechnol. Biofuels.* 8 (2015) 7.
35 doi:10.1186/s13068-014-0187-6.

- [33] V. Puchart, J.W. Agger, J.G. Berrin, A. Várnai, B. Westereng, P. Biely, Comparison of fungal carbohydrate esterases of family CE16 on artificial and natural substrates, *J. Biotechnol.* 233 (2016) 228–236. doi:10.1016/j.jbiotec.2016.07.003.
- [34] G. Mai-Gisondi, O. Turunen, O. Pastinen, N. Pahimanolis, E.R. Master, Enhancement of acetyl xylan esterase activity on cellulose acetate through fusion to a family 3 cellulose binding module, *Enzyme Microb. Technol.* 79–80 (2015) 27–33. doi:10.1016/j.enzmictec.2015.07.001.
- [35] F. Møllerup, E. Master, Influence of a family 29 carbohydrate binding module on the recombinant production of galactose oxidase in *Pichia pastoris*, *Data Br.* 6 (2016) 176–183. doi:10.1016/j.bbagen.2015.10.023.
- [36] P. Gullón, M.J. González-Muñoz, H. Domínguez, J.C. Parajó, Membrane processing of liquors from *Eucalyptus globulus* autohydrolysis, *J. Food Eng.* 87 (2008) 257–265. doi:10.1016/j.jfoodeng.2007.11.032.
- [37] S. Willför, P. Rehn, A. Sundberg, K. Sundberg, B. Holmbom, Recovery of water-soluble acetylglactoglucosylmannans from mechanical pulp of spruce, *Tappi J.* 2 (2003) 27–32.
- [38] W. Shao, J. Wiegel, Purification and characterization of two thermostable acetyl xylan esterases from *Thermoanaerobacterium* sp. strain JW/SL-YS485, *Appl. Environ. Microbiol.* 61 (1995) 729–733. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=167333&tool=pmcentrez&render_type=abstract.
- [39] M. Mastihubová, P. Biely, Lipase-catalysed preparation of acetates of 4-nitrophenyl β -D-xylopyranoside and their use in kinetic studies of acetyl migration, *Carbohydr. Res.* 339 (2004) 1353–1360. doi:10.1016/j.carres.2004.02.016.
- [40] T. Hannuksela, C.H. Du Penhoat, NMR structural determination of dissolved O-acetylated galactoglucosylmannan isolated from spruce thermomechanical pulp, *Carbohydr. Res.* 339 (2004) 301–312. doi:10.1016/j.carres.2003.10.025.
- [41] S.-L. Chong, L. Virkki, H. Maaheimo, M. Juvonen, M. Derba-Maceluch, S. Koutaniemi, M. Roach, B. Sundberg, P. Tuomainen, E.J. Mellerowicz, M. Tenkanen, O-acetylation of glucuronoxylan in *Arabidopsis thaliana* wild type and its change in xylan biosynthesis mutants, *Glycobiology.* 24 (2014) 494–506. doi:10.1093/glycob/cwu017.
- [42] V. Puchart, J.-G. Berrin, M. Haon, P. Biely, A unique CE16 acetyl esterase from *Podospora anserina* active on polymeric xylan, *Appl. Microbiol. Biotechnol.* 99 (2015) 10515–10526. doi:10.1007/s00253-015-6934-1.
- [43] P. Biely, M. Czigárová, J.W. Agger, X.L. Li, V. Puchart, M. Vršanská, V.G.H. Eijssink, B. Westereng, *Trichoderma reesei* CE16 acetyl esterase and its role in enzymatic degradation of

1 acetylated hemicellulose, *Biochim. Biophys. Acta.* 1840 (2014) 516–525.

2 doi:10.1016/j.bbagen.2013.10.008.

- 3 [44] J. Lundqvist, A. Teleman, L. Junel, G. Zacchi, O. Dahlman, F. Tjerneld, H. Stålbrand,
4 Isolation and characterization of galactoglucomannan from spruce (*Picea abies*), *Carbohydr.*
5 *Polym.* 48 (2002) 29–39. doi:10.1016/S0144-8617(01)00210-7.

- 6 [45] E. Margolles-Clark, M. Tenkanen, H. Söderlund, M. Penttilä, Acetyl Xylan Esterase from
7 *Trichoderma reesei* Contains an Active-Site Serine Residue and a Cellulose-Binding
8 Domain, *Eur. J. Biochem.* 237 (1996) 553–560.

9 <http://onlinelibrary.wiley.com/doi/10.1111/j.1432-1033.1996.0553p.x/full> (accessed June 22,
10 2013).

TABLES

Table 1. Specific activity ($\mu\text{mol}/\text{min}/\text{mg}$) of *AnAcXE* (CE1), *OsAcXE* (CE6) and *MtAcE* (CE16) on 4-MUA at pH 6.0 and 7.0.

Enzyme	pH 6.0	pH 7.0
<i>AnAcXE</i> (CE1)	24 ± 0	26 ± 1
<i>OsAcXE</i> (CE6)	201 ± 13	237 ± 5
<i>MtAcE</i> (CE16)	506 ± 7	547 ± 28

n=3; errors indicate standard deviation

Table 2. HSQC NMR analysis of enzyme treated acetyl-4-*O*-methylglucuronoxylan. Mole percent remaining (compared to untreated MGX) of specific acetyl group positions following enzyme treatment.

acetyl-4- <i>O</i> -methylglucuronoxylan (MGX)	Untreated ^a	<i>AnAcXE</i> (CE1)	<i>OsAcXE</i> (CE6)	<i>MtAcE</i> (CE16)
2- <i>O</i> -acetyl-Xylp	24.1 ± 0.2	0.2	1.1	1.1 ± 0.4
3- <i>O</i> -acetyl-Xylp	57.6 ± 0.4	2.3	4.9	2.7 ± 1.4
2,3- <i>O</i> -acetyl-Xylp	6.5 ± 0.3	0.2	1.2	0.2 ± 0.1
(2- <i>O</i> -MeGlc _p A)3- <i>O</i> -acetyl-Xylp	11.7 ± 0.1	8.1	9.0	7.6 ± 0.9
total acetylation (%)	100.0	10.8	16.3	11.6 ± 0.2
reduction acetylation (%)		89.2	83.7	88.4

a. The mole percent of acetyl group positions in the original MGX. n=2 for untreated and *MtAcE* (CE16), otherwise n=1.

Table 3. Specific activity ($\mu\text{mol}/\text{min}/\text{mg}$) of *AnAcXE* (CE1), *OsAcXE* (CE6) and *MtAcE* (CE16) on MGX and GGM. Reactions were performed at pH 7.0 and 40° C; acetyl group release was measured using the acetic acid kit.

Substrate	<i>AnAcXE</i> (CE1)	<i>OsAcXE</i> (CE6)	<i>MtAcE</i> (CE16)
Eucalyptus acetyl-4- <i>O</i> -methylglucuronoxylan (MGX)	5.4 ± 0.7	22.8 ± 3	20.1 ± 2
Spruce acetyl- galactoglucomannan (GGM)	1.2 ± 0.1	1.6 ± 0.3	120 ± 20

n=3 (minimum); errors indicate standard deviation

Table 4. Relative activity (%) towards specific acetyl group positions in eucalyptus acetyl-4-*O*-methylglucuronoxylan (MGX) determined using real time ^1H NMR.

Enzyme	2- <i>O</i> - acetyl-Xylp	3- <i>O</i> - acetyl-Xylp	2,3- <i>O</i> - acetyl-Xylp	(2- <i>O</i> -MeGlc pA)3- <i>O</i> -acetyl-Xylp
<i>AnAcXE</i> (CE1)	32	13	55	ND
<i>OsAcXE</i> (CE6)	40	15	46	ND
<i>MtAcE</i> (CE16)	45	ND	55	ND

Relative activity was calculated as follows: (slope/mgenzyme/(% acetyl groups at the specific position in untreated MGX)/ $\sum$ (slope/mgenzyme/(% acetyl groups at the specific position in untreated MGX) *100. ND, not determined given that corresponding slope values were insignificantly different from zero (Suppl. Fig. 4).

Table 5. Relative activity (%) towards specific acetyl group positions in spruce acetyl-galactoglucomannan (GGM) determined using real time ^1H NMR.

Enzyme	2- <i>O</i> - acetyl-Man p	3- <i>O</i> - acetyl-Man p	2- <i>O</i> -acetyl- Man p ; H2	3- <i>O</i> -acetyl- Man p ; H3
<i>AnAcXE</i> (CE1)	39	61	31	69
<i>OsAcXE</i> (CE6)	56	44	83	17
<i>MtAcE</i> (CE16)	50	50	84	16

Calculation for relative activity: $(\text{slope}/\text{mg}_{\text{enzyme}}/(\% \text{ acetyl groups at the specific position in untreated GGM}) / \sum (\text{slope}/\text{mg}_{\text{enzyme}}/(\% \text{ acetyl groups at the specific position in untreated GGM}))$

*100.

FIGURE LEGENDS

Fig. 1. Molecular structures of acetyl-4-*O*-methylglucuronoxylan (MGX) (A) and acetyl-galactoglucomannan (GGM) (B). Acetyl group positions are shown in red.

Fig. 2. ¹H NMR spectra following enzyme action on acetyl-4-*O*-methylglucuronoxylan (MGX) and acetyl-galactoglucomannan (GGM) in buffered D₂O at 40° C. The first spectrum was collected after 10 min of reaction; subsequent spectra were collected every 1.8 min. (A) MGX before and after treatment with *AnAcXE* (CE1) (5 µg/mg MGX), *OsAcXE* (CE6) (1 µg/mg MGX), *MtAcE* (CE16) (1 µg/mg MGX); (B) GGM before and after treatment with *AnAcXE* (CE1) (10 µg/mg GGM), *OsAcXE* (CE6) (10 µg/mg GGM), *MtAcE* (CE16) (1 µg/mg GGM).

Fig. 3. Enzyme stability at pH 6.0 and 40°C for up to 24 h. Relative activity was measured using 4-MUA. Average values are indicated above each bar. n=3; error bars indicate standard deviation.

Fig. 4. Deacetylation of acetyl-4-*O*-methylglucuronoxylan (MGX; black bars) and acetyl-galactoglucomannan (GGM; grey bars). Percent of acetyl group content released by *AnAcXE* (CE1), *OsAcXE* (CE6), and *MtAcE* (CE16) (10 µg enzyme /1 mg substrate) measured after 24 h at 40°C and pH 6.0, using the Acetic Acid kit from Megazyme. Average values are indicated above each bar. Non-enzymatic release of acetyl groups from MGX and GGM at pH 6.0 after 24 h was 4 % and 2%, respectively. n=3; error bars represent standard deviation.

Fig. 5. HSQC spectra of GGM in DMSO at 60° C. Acetyl-galactoglucomannan (GGM) in DMSO-d₆ before and after treatment with *AnAcXE* (CE1), *OsAcXE* (CE6), *MtAcE* (CE16) (10 µg enzyme /1 mg substrate) for 24 h at pH 6.0 and 40°C. Scaling for the native substrate was adjusted to show all available peaks. Acetylated O-2 (H2) and O-3 (H3) positions in the anomeric region are shown due to the partial overlap of signals in the acetyl group region between 2.5-2.0 ppm.

A.

B.

Fig. 1.

Fig. 2.

Fig. 3.

Fig. 4

Fig. 5.

SUPPLEMENTAL TABLE

Suppl.Table 1. Reported compositions (mol%) of acetyl-4-*O*-methylglucuronoxylan (MGX), acetyl-galactoglucomannan (GGM), determined using NMR or by acid methanolysis/gas-chromatography (Meth-GC)

Sample	Extraction/ Determination method	β -D- Xylp	Me-GlcpA	β -D-Manp	β -D-Glcp	α -D-Galp	O-Ac (DS)	Ref.
MGX (Birch)	DMSO/ NMR	68	3	-	-	-	29 (0.4)	[9]
MGX (Beech)		70	4	-	-	-	26 (0.3)	
MGX (Aspen)	Microwave/ NMR	60	6	-	-	-	35 (0.6)	[7]
GGM (Birch)	Hot water/ NMR	-	-	68	32	-	(0.2)	[10]
GGM (Spruce)		2-9	1-2	43-65	7-19	7-20	(0.5)	[3]
TMP or wood	Meth-GC							

DS: degree of substitution

- not determined or not detected

TMP: thermomechanical pulp

SUPPLEMENTAL FIGURES

Suppl.Fig. 1. SDS-PAGE analysis of purified enzymes (15 µg). Lanes, PageRuler™ Prestained Protein Ladder, 10 to 180 kDa (ThermoFisher Scientific; USA), 2: *AnAcXE* (CE1); 3: *OsAcXE* (CE6); 4: *MtAE* (CE16).

Suppl.Fig. 2. Optimum pH of selected enzymes measured using 4-MUA. (A) *AnAcXE*(CE1); (B) *OsAcXE* (CE6), and (C) *MtAcE* (CE16).

Suppl.Fig. 3. Optimum pH of (A) *AnAcXE* (CE1) and (B) *MtAcE* (CE16) using MGX.

Suppl.Fig. 4. Normalized resonance signals from H¹ NMR plotted against time. (A) Plots showing activity of each enzyme towards each acetyl group in MGX and GGM, (B) Slopes calculated from plots shown in (A). Prob(F) values were obtained using ANOVA (Origin 2017 software); Prob(F) values >0.05 indicate insignificant difference from zero.

Suppl.Fig. 1

A. *AnAcXE* (CE1)

B. *OsAcXE* (CE6)

C. *MtAcE* (CE16)

Suppl.Fig. 2

A. *AnAcXE* (CE1)

B. *MtAcE* (CE16)

Suppl.Fig. 3

Suppl.Fig. 4A

Calculated slopes derived from enzyme-treated MGX								
Enzyme	2- <i>O</i> - acetyl-Xylp	Prob (F) 2- <i>O</i> - acetyl-Xylp	3- <i>O</i> - acetyl-Xylp	Prob (F) 3- <i>O</i> - acetyl-Xylp	2,3- <i>O</i> - acetyl-Xylp	Prob (F) 2,3- <i>O</i> - acetyl-Xylp	(2- <i>O</i> - MeGlc <p>A</p>) 3- <i>O</i> -acetyl- Xylp	Prob (F) (2- <i>O</i> - MeGlc <p>A</p>) 3- <i>O</i> -acetyl- Xylp
<i>AnAcXE</i> (CE1)	0.11 ± 0.02	3.5 x 10 ⁻⁶	0.06 ± 0.01	7.8 x 10 ⁻⁴	0.10 ± 0.002	0.0	0.002 ± 0.002	3.1 x 10 ⁻¹
<i>OsAcXE</i> (CE6)	0.11 ± 0.01	1.9 x 10 ⁻¹¹	0.05 ± 0.01	3.9 x 10 ⁻⁷	0.07 ± 0.003	2.1 x 10 ⁻¹²	0.001 ± 0.002	6.1 x 10 ⁻¹
<i>MtAcE</i> (CE16)	0.11 ± 0.01	4.9 x 10 ⁻⁷	0.003 ± 0.007	7.1 x 10 ⁻¹	0.07 ± 0.01	1.3 x 10 ⁻⁸	0.01 ± 0.003	6.3 x 10 ⁻²
Calculated slopes derived from enzyme-treated GGM								
Enzyme	2- <i>O</i> - acetyl-Manp	Prob (F) 2- <i>O</i> - acetyl-Manp	3- <i>O</i> - acetyl-Manp	Prob (F) 3- <i>O</i> - acetyl-Manp	2- <i>O</i> - acetyl-Manp; H2	Prob (F) 2- <i>O</i> - acetyl-Manp; H2	3- <i>O</i> - acetyl-Manp; H3	Prob (F) 3- <i>O</i> - acetyl-Manp; H3
<i>AnAcXE</i> (CE1)	0.25 ± 0.02	1.1 x 10 ⁻⁹	0.26 ± 0.02	5.3 x 10 ⁻⁹	0.08 ± 0.02	5.2 x 10 ⁻⁴	0.24 ± 0.03	8.1 x 10 ⁻⁸
<i>OsAcXE</i> (CE6)	0.23 ± 0.03	3.9 x 10 ⁻⁷	0.13 ± 0.02	1.1 x 10 ⁻⁶	0.19 ± 0.01	2.3 x 10 ⁻¹⁰	0.05 ± 0.01	7.7 x 10 ⁻⁵
<i>MtAcE</i> (CE16)	0.31 ± 0.01	2.2 x 10 ⁻¹⁶	0.19 ± 0.01	3.1 x 10 ¹³	0.36 ± 0.01	0	0.09 ± 0.01	6.3 x 10 ⁻⁸

Suppl.Fig. 4B