This is an electronic reprint of the original article.
This reprint may differ from the original in pagination and typographic detail.

Volovik, Grigory

Varying Newton constant and black hole to white hole quantum tunneling

Published in:
Universe

DOI:
[10.3390/UNIVERSE6090133](https://doi.org/10.3390/UNIVERSE6090133)

Published: 01/01/2020

Document Version
Publisher's PDF, also known as Version of record

Published under the following license:
CC BY

Please cite the original version:
Volovik, G. (2020). Varying Newton constant and black hole to white hole quantum tunneling. *Universe*, 6(9), 1-8. [133]. <https://doi.org/10.3390/UNIVERSE6090133>

Article

Varying Newton Constant and Black Hole to White Hole Quantum Tunneling

Grigory Volovik ^{1,2}
¹ Low Temperature Laboratory, Aalto University, P.O. Box 15100, FI-00076 Aalto, Finland; grigori.volovik@aalto.fi

² Landau Institute for Theoretical Physics, acad. Semyonov av., 1a, 142432 Chernogolovka, Russia

Received: 31 July 2020; Accepted: 22 August 2020; Published: 23 August 2020

Abstract: The thermodynamics of black holes is discussed for the case, when the Newton constant G is not a constant, but it is the thermodynamic variable. This gives for the first law of the Schwarzschild black hole thermodynamics: $dS_{\text{BH}} = -AdK + \frac{dM}{T_{\text{BH}}}$, where the gravitational coupling $K = 1/4G$, M is the black hole mass, A is the area of horizon, and T_{BH} is Hawking temperature. From this first law, it follows that the dimensionless quantity M^2/K is the adiabatic invariant, which, in principle, can be quantized if to follow the Bekenstein conjecture. From the Euclidean action for the black hole it follows that K and A serve as dynamically conjugate variables. Using the Painlevé–Gullstrand metric, which in condensed matter is known as acoustic metric, we calculate the quantum tunneling from the black hole to the white hole. The obtained tunneling exponent suggests that the temperature and entropy of the white hole are negative.

Keywords: black hole; white hole; quantum tunneling; negative entropy

1. Introduction

Bekenstein [1] proposed that the horizon area A is an adiabatic invariant and, thus, can be quantized according to the Ehrenfest principle that classical adiabatic invariants may correspond to observables with discrete spectrum. Here, we consider the black hole thermodynamics in case when the gravitational coupling K is the variable thermodynamic quantity. The variable K modifies the first law of the black hole thermodynamics and leads to the alternative adiabatic invariant, which is dimensionless and, thus, in principle, can be quantized.

On the quantum level, the gravitational coupling K becomes the variable, which is dynamically conjugate to the black hole area A . This allows us to study the black hole to white hole transition while using the semiclassical description of the quantum tunneling and considering the trajectory in the (K, A) phase space, which connects the black and white holes. We also consider the temperature and entropy of the white hole, which is formed from the black hole by quantum tunneling and find that both the temperature and entropy of the white hole are negative.

2. Modified First Law of Black Hole Thermodynamics

The Einstein–Hilbert gravitational action is

$$S_{\text{grav}} = \frac{1}{4\pi} \int d^3x dt \sqrt{-g} K \mathcal{R}, \quad (1)$$

where $\mathcal{R}$ is the scalar curvature, and we choose the gravitational coupling $K = 1/(4G) = 1/(4l_{\text{Planck}}^2)$. In the modified gravity theories, such as the scalar-tensor and $f(R)$ theories (see e.g., [2] and the latest paper [3] with references therein), the effective Newton “constant” G can be space-time dependent

and, thus, is not fundamental. Additionally, it was suggested that there is connection between the gravitational coupling K and the fine structure constant, see Refs. [4–10]. In the chiral superfluid $^3\text{He-A}$ the coupling K in the effective gravity is determined by physics on microscopic (atomic) level [11], and it depends on coordinates and temperature.

Here, we assume that the variables, which enter the Einstein action—the scalar Riemann curvature R and the gravitational coupling constant K —are the local thermodynamic variables, which are similar to temperature, pressure, chemical potential, number density, etc. The Riemann curvature as the covariant quantity may serve as one of the thermodynamical characteristics of the macroscopic matter [12]. If so, the gravitational coupling constant K in front of the scalar curvature in Einstein action also becomes the thermodynamic quantity, see Ref. [13]. The application of this thermodynamics to the global object, such as black hole, suggests that the variable K (actually its asymptotic value at infinity) should enter the thermodynamic laws for the black hole. The corresponding thermodynamical conjugate to the global K is obtained by the volume integral of the Riemann curvature.

In terms of this coupling K , the Hawking temperature of Schwarzschild black hole and its Bekenstein entropy are:

$$T_{\text{BH}} = \frac{K}{2\pi M}, \quad S_{\text{BH}} = \frac{\pi M^2}{K}. \quad (2)$$

Then, using the black hole area $A = \pi M^2/K^2$, the gravitational coupling $K = 1/4G$, the Hawking temperature $T_{\text{BH}} = M/2AK = K/2\pi M$, and the black hole entropy $S_{\text{BH}} = AK$, one obtains:

$$\begin{aligned} dS_{\text{BH}} &= d(AK) = \pi d(M^2/K) = \\ &= -\pi \frac{M^2}{K^2} dK + 2\pi \frac{M}{K} dM. \end{aligned} \quad (3)$$

This suggests the following modification of the first law of black hole thermodynamics in case if K is a thermodynamic variable:

$$dS_{\text{BH}} = -AdK + \frac{dM}{T_{\text{BH}}}. \quad (4)$$

This modification is similar to the modification in terms of the moduli fields [14]. However, in our case, the thermodynamic variable, which is conjugate to the thermodynamic variable K , is the product of the black hole area and the black hole temperature, AT_{BH} . On the other hand, in dynamics, K and A are canonically conjugate, see Section 4.

In general, the variable K is local and depends on space coordinate, but in the same way as for the moduli fields [14] the black hole thermodynamics is determined by the asymptotic value of K at spatial infinity. In Equation (4), $K \equiv K(\infty)$ is the global quantity, which characterizes the quantum vacuum in full equilibrium, i.e., far from the black hole. The variable K allows for us to study the transition to the vacuum without gravity, i.e., to the vacuum where $K \rightarrow \infty$ and, thus, $G \rightarrow 0$, see Section 4.

3. Adiabatic Change of K and Adiabatic Invariant

Let us change K and M adiabatically, i.e., at constant entropy. Then the equation $dS_{\text{BH}} = 0$ gives

$$\frac{dM}{dK} = AT_{\text{BH}} = \frac{M}{2K}. \quad (5)$$

This shows that $M^2/K = \text{const}$ is the adiabatic invariant for the spherical neutral black hole and, thus, according to the Bekenstein conjecture [1], it can be quantized in quantum mechanics:

$$\frac{M^2}{K} = aN. \quad (6)$$

Here, N is integer, and a is some fundamental dimensionless parameter of order unity. If this conjecture is correct, for the entropy of Schwarzschild black hole one has

$$S_{\text{BH}}(N) = \pi \frac{M^2}{K} = \pi a N. \quad (7)$$

The Bekenstein idea on the role of adiabatic invariants in quantization of the black hole requires further consideration, see some approaches to that in Refs. [15–20]. In particular, the similarity between the energy levels of Schwarzschild black hole and the hydrogen atom has been suggested [21,22]. We leave this problem for the future. This consideration should be supported by microscopic theory, see e.g., [23]. The so-called q -theory can be exploited, which allows us to consider dark energy, dark matter, black holes, and the varying gravitational coupling in the frame of the same effective theory of the quantum vacuum [13,24,25].

4. A and K as Canonically Conjugate Variables and Black-Hole—White-Hole Quantum Tunneling

Till now, we discussed the black hole thermodynamics, where the adiabatic invariant M^2/K arises. Now, we move to the dynamics of black hole, and apply the varying K approach to the consideration of the quantum tunneling of the black hole to white hole. The quantum tunneling can be considered in a semiclassical approximation in the same manner as Hawking radiation is discussed in terms of semiclassical quantum tunneling [26–29]. For ${}^3\text{He-A}$, the Hawking radiation in terms of quantum tunneling was considered in Ref. [30].

The quantum mechanical treatment of the black hole can be obtained, if one finds the relevant canonically conjugate variables describing the black hole dynamics. Such an approach has been suggested in Ref. [31], where the canonically conjugate variables have been determined in the Euclidean time. Because, in Euclidean time, the action for the black hole is equal to its entropy, $I_E = S_{\text{BH}}$, in the theory with varying gravitational coupling K , the proper canonically conjugate variables are the gravitational coupling K and the black hole area A . This allows us to consider the quantum mechanical tunneling from the black hole to the white hole, which was discussed in Refs. [32–40] and references therein.

The process of quantum tunneling of macroscopic objects is well known in condensed matter physics, where the collective variables are used, which describe the collective dynamics of a macroscopic body [41,42]. In particular, in the quantum tunneling creation of quantized vortices in superfluids [43] and superconductors [44], the dynamically conjugate variables are the area of the vortex ring and its coordinate along the normal to the ring. This approach provides the correct semiclassical tunneling exponent without consideration of the details of the structure of the object on the microscopic level. In the same manner, it looks reasonable that, in case of tunneling from the black hole to the white hole, described by the collective variables K and A , the microscopic processes are not important. Note the difference from consideration in Section 3, where K varies in the adiabatic regime, i.e., at fixed entropy S_{BH} , while the area A , temperature T_{BH} , and mass M follow the variation of K . In the dynamic regime, which is relevant for the description of quantum tunnelling, the parameter K varies at fixed energy (mass M), while the area A , the temperature T_{BH} , and entropy S_{BH} follow the variation of K .

In the case of Hawking radiation, the tunneling of particle from inside to outside the horizon is obtained by evaluating the action of the particle [26,28,30,45,46]. The tunneling rate is determined by imaginary part of the action and can be obtained while using the path in the complex plane. The calculation of the tunneling exponent demonstrates that it is proportional to the change of the black hole entropy after radiation of a particle, $p \propto e^{\Delta S_{\text{BH}}}$, see Refs. [26,45,46]. The same result can also be obtained using the conjugate variables K and A in the Euclidean action, and the path $\int A(K) dK$ at fixed M with real K . The direct connection between the Hawking radiation and black hole tunneling will be discussed in Section 5.

As in the case of the semiclassical consideration of the Hawking radiation in terms of the quantum tunneling [26,28,30], we shall use the Painleve–Gullstrand coordinate system [47,48] with the metric:

$$ds^2 = -dt^2(1 - \mathbf{v}^2) - 2dt \, d\mathbf{r} \cdot \mathbf{v} + d\mathbf{r}^2. \quad (8)$$

Here, the vector $v_i(\mathbf{r}) = g_{0i}(\mathbf{r})$ is the velocity of the free-falling observer, which crosses the horizon. In condensed matter the analog of this metric is the so-called acoustic metric [49]. In superfluids, the velocity v_i is played by superfluid velocity. The analogs of the black hole and white hole horizons described by this metric can be also reproduced in the Dirac and Weyl topological semimetals, where the horizon takes place on the boundary between different types of Dirac or Weyl materials [50–53].

For the Schwarzschild black hole, one has

$$\mathbf{v}(\mathbf{r}) = \mp \hat{\mathbf{r}} \sqrt{\frac{r_H}{r}} = \mp \hat{\mathbf{r}} \sqrt{\frac{M}{2rK}}, \quad (9)$$

where r_H is the radius of the horizon; the minus sign corresponds to the black hole and the plus sign describes the white hole [32]. Note that, in the theory with the variable gravitational coupling, the sign changes at the singularity $K = \infty$ (or at $G = 0$), when the black hole shrinks to a point and then expands as a white hole. In terms of variable K , the point $K = \infty$ serves as the branch point, where the velocity of the freely falling observer changes sign.

It is important that the vector $\mathbf{v}$, which is normal to the surface of the horizon, is the velocity of the free falling observer, who crosses the horizon. For the observer, who crosses the black hole horizon from outside, and for the observer, who crosses the white hole horizon from inside, these two directions are opposite. For these two observers the area of the horizons has different sign. This means that, at the branch point of the trajectory, the horizon area A changes sign: it crosses zero at $K = \infty$ and becomes negative on the white-hole side of the process, $A \rightarrow -A$. This could also mean that, due to connection between the area and entropy, the white hole may have negative entropy, which we discuss in Section 5.

The quantum tunneling exponent is usually determined by the imaginary part of the action on the trajectory, which transforms the black hole to white hole. In terms of Euclidean action, one has:

$$p \propto \exp(-I_{\text{BH} \rightarrow \text{WH}}), \quad I_{\text{BH} \rightarrow \text{WH}} = \int_C A(K') dK'. \quad (10)$$

Here, the semiclassical trajectory C is at $M = \text{const}$, and thus $A(K') = \pm \pi M^2 / (K')^2$. Along this trajectory the variable K' changes from K to the branch point at $K' = \infty$, and then from $K' = \infty$ to $K' = K$ along the other branch, where the area $A(K') < 0$. The integral gives the tunneling exponent of the transition to the white hole

$$I_{\text{BH} \rightarrow \text{WH}} = 2\pi M^2 \int_K^\infty \frac{dK'}{K'^2} = 2\pi \frac{M^2}{K}, \quad (11)$$

and the transition probability is:

$$p \propto \exp\left(-2\pi M^2 / K\right) = \exp(-2S_{\text{BH}}). \quad (12)$$

5. White Hole Entropy and Temperature

The result (12) can be also obtained using the Hawking radiation from the black hole, where the tunneling exponent is proportional to $e^{\Delta S_{\text{BH}}}$, see Refs. [26,45,46]. Let us consider the process, in which the particle escapes the black hole by quantum tunneling and then it tunnels to the white hole through the white hole horizon. This process occurs at the fixed total mass M . The tunneling exponent for this

process to occur is $e^{(\Delta S_{\text{BH}} + \Delta S_{\text{WH}})}$. The summation of all the processes of the tunneling of matter from the black hole to the formed white hole finally gives Equation (12):

$$p \propto e^{\sum(\Delta S_{\text{BH}} + \Delta S_{\text{WH}})} = e^{2\sum\Delta S_{\text{BH}}} = \exp(-2S_{\text{BH}}). \quad (13)$$

Here, we took into account the (anti)symmetry in the dynamics of black and white holes in the process of quantum tunneling, $\sum\Delta S_{\text{BH}} = \sum\Delta S_{\text{WH}}$.

As in the case of quantum tunneling in the Hawking radiation process, the probability in Equation (12) has the thermodynamic meaning as thermodynamic fluctuation $e^{\Delta S}$ [54]. In our case, the total change of the entropy in the process of the tunneling from black to white hole is $\Delta S = S_{\text{WH}} - S_{\text{BH}}$. According to Equation (13), this change is equal to $-2S_{\text{BH}}$. As a result, one obtains that the entropy of the white hole is equal with the opposite sign to the entropy of the black hole with the same mass

$$S_{\text{WH}}(M) = -S_{\text{BH}}(M). \quad (14)$$

This means that the white hole, which is obtained by quantum tunneling from the black hole and, thus, has the same mass M as the black hole, has the negative temperature $T_{\text{WH}} = -T_{\text{BH}}$ and the negative area $A_{\text{WH}} = -A_{\text{BH}}$, which together produce the negative entropy $S_{\text{WH}} = -S_{\text{BH}}$.

The negative temperature is a well defined quantity in condensed matter. It typically takes place in the subsystem of nuclear spins, where the energy is restricted from above. Different thermodynamic phase transitions occurring at $T < 0$ have been experimentally observed in magnetic systems, see e.g., Ref. [55]. The negative temperature states are unstable both before and after the magnetic phase transition. In principle, the state with $T < 0$ is hotter than the state $T > 0$, since the heat flows from the negative- to the positive-temperature system. If the black hole and white hole are in some contact, then the heat will escape from the white hole and be absorbed by the black hole.

The transition from positive to negative T occurs on the path $A(K)dK$ via the point $K = \infty$, where $T(K) = \infty$. The transition via infinite temperature is the analytic route to the thermodynamically unstable states with negative T , see e.g., Refs. [56,57], where the transition to anti-spacetime [58–60] has been considered. In spin systems, the $T < 0$ state is typically obtained by reversing the magnetic field, which looks like crossing $T = 0$.

In the black hole physics, the negative temperature has been discussed for the inner horizon of the Kerr and charged black holes, see Ref. [61] and the references therein. The entropy of the inner horizon has been considered as positive. However, the arguments in Ref. [61] do not exclude the possibility of the opposite situation, when the temperature of the inner horizon is positive, while the entropy is negative: the equation $T_+S_+ + T_-S_- = 0$ remains valid.

The discussed transition from the black hole to the white hole with the same mass M is not the thermodynamic transition. It is the quantum process of tunneling between the two quantum states. It is one of many routes of the black hole evaporation, including the formation of small white hole on the late stage of the decay [32–40]. The uniqueness of this route together with hidden information and (anti)symmetry between the black and white holes, is the possible origin of the negative entropy of the white hole.

6. Discussion

We considered the dynamics and thermodynamics of the black hole in the case of the varying gravitational coupling K . The gravitational coupling K serves as the thermodynamic variable, which is thermodynamic conjugate to $A_{\text{BH}}T_{\text{BH}}$, where A_{BH} and T_{BH} are correspondingly the area of the black hole horizon and Hawking temperature. The corresponding first law of the black hole is modified, see Equation (4). The corresponding adiabatic invariant is the entropy $S_{\text{BH}} = KA_{\text{BH}}$, and it is this invariant that should be quantized, if the Bekenstein conjecture is correct. This is in agreement with observation of Ted Jacobson [62], that it is the entropy that does not change under renormalization of K , rather than the area. This suggests the alternative quantization scheme for the black hole. While K

and A_{BH} are dimensionful and cannot be quantized, the entropy is dimensionless and, thus, can be quantized in terms of some fundamental numbers.

On the quantum level, the dynamically conjugate variables of the black hole physics are K and A_{BH} . This allows us to consider the transition from the black hole to the white hole as quantum tunneling in the semiclassical approximation, which is valid when the action is large. The classical trajectory of the black hole crosses the branch point at $K = \infty$, and then continues along the other branch, where the area $A(K) < 0$, which corresponds to the white hole. The obtained tunneling exponent $\exp(-2S_{\text{BH}})$ demonstrates that the transition can be considered as thermodynamic fluctuation, if the entropy of the white hole (with the same mass M as the black hole) is negative, $S_{\text{WH}} = -S_{\text{BH}}$. The latter also suggests that this white hole has negative temperature $T_{\text{WH}} = -T_{\text{BH}}$.

Funding: This work has been supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant Agreement No. 694248).

Conflicts of Interest: The author declares no conflict of interest

Acknowledgments: I thank J. Barrow and A. Starobinsky for criticism and H. Terazawa for correspondence.

References

1. Bekenstein, J. D. The quantum mass spectrum of the Kerr black hole, *Lett. Nuovo Cimento* **1974**, *11*, 467.
2. Starobinsky, A.A. A new type of isotropic cosmological models without singularity. *Phys. Lett. B* **1980**, *9*, 99. [[CrossRef](#)]
3. Barrow, J.D. Finite action principle revisited. *Phys. Rev. D* **2020**, *101*, 023527. [[CrossRef](#)]
4. Landau, L.D. *Niels Bohr and the Development of Physics*; Pauli, W., Rosenfeld, L., Weisskopf, V., Eds.; McGraw-Hill: New York, NY, USA, 1955; p. 52.
5. Akama, K. Pregeometry. In *Gauge Theory and Gravitation*; Nakanishi, N., Nariai, H., Eds.; Springer: Berlin, Germany, 1983; pp. 267–271.
6. Akama, K.; Chikashige, Y.; Matsuki, T.; Terazawa, H. Gravity and electromagnetism as collective phenomena: A derivation of Einstein's general relativity. *Prog. Theor. Phys* **1978**, *60*, 868. [[CrossRef](#)]
7. Terazawa, H. High energy physics in the 21st century. In Proceedings of the 22nd International Workshop on the Fundamental Problems of High Energy Physics and Field Theory, Protvino, Moscow Region, Russian Federation, 23–25 June 1999; KEK Preprint 99-46, July 1999.
8. Terazawa, H.; Chikashige, Y.; Akama, K.; Matsuki, T. Simple relation between the fine-structure and gravitational constants. *Phys. Rev. D* **1977**, *15*, 1181–1183. [[CrossRef](#)]
9. Terazawa, H. Cosmological origin of mass scales. *Phys. Lett. B* **1981**, *101*, 43. [[CrossRef](#)]
10. Klinkhamer, F.R.; Volovik, G.E. Merging gauge coupling constants without Grand Unification. *J. Experiment. Theor. Phys. Lett.* **2005**, *81*, 551–555. [[CrossRef](#)]
11. Volovik, G.E. *The Universe in a Helium Droplet*; Clarendon Press: Oxford, UK, 2003.
12. Pronin, P.; Kulikov, I. Local quantum statistics in arbitrary curved space-time. *Pramana* **1987**, *28*, 355–359. [[CrossRef](#)]
13. Klinkhamer, F.R.; Volovik, G.E. $f(R)$ cosmology from q -theory. *JETP Lett.* **2008**, *88*, 289–294. [[CrossRef](#)]
14. Gibbons, G.; Kallosh, R.; Kol, B. Moduli, scalar charges, and the first law of black hole thermodynamics. *Phys. Rev. Lett.* **1996**, *77*, 4992. [[CrossRef](#)]
15. Horowitz, G.T.; Strominger, A. Counting states of near extremal black holes. *Phys. Rev. Lett.* **1996**, *77*, 2368. [[CrossRef](#)] [[PubMed](#)]
16. Barvinsky, A.; Das, S.; Kunstatter, G. Quantum mechanics of charged black holes. *Phys. Lett. B* **2001**, *517*, 415–420. [[CrossRef](#)]
17. Barvinsky, A.; Das, S.; Kunstatter, G. Discrete spectra of charged black holes. *Foundat. Phys.* **2002**, *32*, 1851–1862. [[CrossRef](#)]
18. Ansorg, M.; Hennig, J. The inner Cauchy horizon of axisymmetric and stationary black holes with surrounding matter. *Class. Quantum Grav.* **2008**, *25*, 22. [[CrossRef](#)]
19. Visser, M. Quantization of area for event and Cauchy horizons of the Kerr-Newman black hole. *J. High Energy Phys.* **2012**, *2012*, 23. [[CrossRef](#)]

20. Tharanath, R.; Kuriakose, V.C. Thermodynamics and spectroscopy of Schwarzschild black hole surrounded by Quintessence. *Mod. Phys. Lett. A* **2013**, *28*, 1350003. [[CrossRef](#)]
21. Bekenstein, J.D. Quantum black holes as atoms. In Proceedings of the Eight Marcel Grossmann Meeting, Jerusalem, Israel, 22–27 June 1997; Piran, T., Ruffini, R., Eds.; World Scientific: Singapore, 1999; pp. 92–111.
22. Corda, C. Time dependent Schrödinger equation for black hole evaporation: No information loss. *Ann. Phys.* **2015**, *353*, 71–72. [[CrossRef](#)]
23. Carlip, S. Black hole thermodynamics. *Int. J. Mod. Phys. D* **2014**, *23*, 1430023. [[CrossRef](#)]
24. Klinkhamer, F.R.; Volovik, G.E. Dark matter from dark energy in q -theory. *JETP Lett.* **2017**, *105*, 74–77. [[CrossRef](#)]
25. Klinkhamer, F.R.; Santillan, O.; Volovik, G.E.; Zhou, A. Vacuum energy decay from a q -bubble. *Physics* **2019**, *1*, 321–338. [[CrossRef](#)]
26. Parikh, M.K.; Wilczek, F. Hawking radiation as tunneling. *Phys. Rev. Lett.* **2000**, *85*, 5042. [[CrossRef](#)] [[PubMed](#)]
27. E.T. Akhmedov, V. Akhmedova, D. Singleton, Hawking temperature in the tunneling picture, *Phys. Lett. B* **2006**, *642*, 124–128.
28. Volovik, G.E. Particle decay in de Sitter spacetime via quantum tunneling. *JETP Lett.* **2009**, *90*, 1–4. [[CrossRef](#)]
29. Vanzo, L.; Acquaviva, G.; Di Criscienzo, R. Tunnelling methods and Hawking’s radiation: Achievements and prospects. *Class. Quantum Grav.* **2011**, *28*, 183001. [[CrossRef](#)]
30. Volovik, G.E. Simulation of Painleve-Gullstrand black hole in thin $^3\text{He-A}$ film. *JETP Lett.* **1999**, *69*, 705–713. [[CrossRef](#)]
31. Majhi, B.R.; Vagenas, E.C. Black hole spectroscopy via adiabatic invariance. *Phys. Lett. B* **2011**, *701*, 623–625. [[CrossRef](#)]
32. Barcelo, C.; Carballo-Rubio, R.; Garay, L.J. Mutiny at the white-hole district. *Int. J. Mod. Phys. D* **2014**, *23*, 1442022. [[CrossRef](#)]
33. Barcelo, C.; Carballo-Rubio, R.; Garay, L.J. Exponential fading to white of black holes in quantum gravity. *Class. Quantum Grav.* **2017**, *34*, 105007. [[CrossRef](#)]
34. Bianchi, E.; Christodoulou, M.; D’Ambrosio, F.; Haggard, H.M.; Rovelli, C. White holes as remnants: A surprising scenario for the end of a black hole. *Class. Quantum Grav.* **2018**, *35*, 225003. [[CrossRef](#)]
35. Martin-Dussaud, P.; Rovelli, C. Evaporating black-to-white hole. *Class. Quantum Grav.* **2019**, *36*, 245002. [[CrossRef](#)]
36. Rovelli, C. Viewpoint: Black hole evolution traced out with loop quantum gravity. *Physics* **2018**, *11*, 127. [[CrossRef](#)]
37. Achour, J.B.; Uzan, J.P. Effective theory of a pulsating Planck star. *arXiv* **2020**, arXiv:2001.06153.
38. Achour, J.B.; Brahma, S.; Uzan, J.P. Bouncing compact objects. Part I. Quantum extension of the Oppenheimer-Snyder collapse. *J. Cosmol. Astropart. Phys.* **2020**, *03*, 041. [[CrossRef](#)]
39. Achour, J.B.; Brahma, S.; Mukohyama, S.; Uzan, J.P. Consistent black-to-white hole bounces from matter collapse. *arXiv* **2020**, arXiv:2004.1297.
40. Bodendorfer, N.; Mele, F.M.; Münch, J. Mass and horizon Dirac observables in effective models of quantum black-to-white hole transition. *arXiv* **2019**, arXiv:1912.00774.
41. Lifshitz, I.M.; Kagan, Y. Quantum kinetics of phase transitions at temperatures close to absolute zero. *JETP* **1972**, *35*, 206–214.
42. Iordanskii, S.V.; Finkel’shtein, A.M. Effect of quantum fluctuations on the lifetimes of metastable states in solids. *JETP* **1972**, *35*, 215.
43. Volovik, G.E. Quantum mechanical creation of vortices in superfluids. *JETP Lett.* **1972**, *15*, 81.
44. Blatter, G.; Feigel’man, M.V.; Geshkenbein, V.B.; Larkin, A.I.; Vinokur, V.M. Vortices in high-temperature superconductors. *Rev. Mod. Phys.* **1994**, *66*, 1125–1388. [[CrossRef](#)]
45. Keski-Vakkuri, E.; Kraus, P. Microcanonical D-branes and back reaction. *Nucl. Phys. B* **1997**, *491*, 249–262. [[CrossRef](#)]
46. Berezin, V.A.; Boyarsky, A.M.; Neronov, A.Y. On the mechanism of Hawking radiation. *Gravit. & Cosmol.* **1999**, *5*, 16–22.
47. Painlevé, P. La mécanique classique et la théorie de la relativité. *C. R. Acad. Sci. (Paris)* **1921**, *173*, 677.
48. Gullstrand, A. Allgemeine Lösung des statischen Einkörper-problems in der Einsteinschen Gravitations-theorie. *Arkiv. Mat. Astron. Fys.* **1992**, *16*, 8.

49. Unruh, W. Experimental black hole evaporation. *Phys. Rev. Lett.* **1981**, *46*, 1351–1353. [[CrossRef](#)]
50. Volovik, G.E. Black hole and Hawking radiation by type-II Weyl fermions. *JETP Lett.* **2016**, *104*, 645–648. [[CrossRef](#)]
51. Volovik, G.E.; Zhang, K. Lifshitz transitions, type-II Dirac and Weyl fermions, event horizon and all that. *J. Low Temp. Phys.* **2017**, *189*, 276–299. [[CrossRef](#)]
52. Kedem, Y.; Bergholtz, E.J.; Wilczek, F. Black and white holes at material junctions. *arXiv* **2020**, arXiv:2001.02625.
53. Nissinen, J.; Volovik, G.E. Type-III and IV interacting Weyl points. *JETP Lett.* **2017**, *105*, 447–452. [[CrossRef](#)]
54. Landau, L.D.; Lifshitz, E.M. Course of Theoretical Physics. In *Statistical Physics*; Pergamon Press: Oxford, UK, 1980; Volume 5.
55. Hakonen, P.J.; Vuorinen, R.T. Nuclear ferromagnetic ordering in silver at negative nanokelvin temperatures. *J. Low Temp. Phys.* **1992**, *89*, 177–186. [[CrossRef](#)]
56. Volovik, G.E. Negative temperature for negative lapse function. *JETP Lett.* **2019**, *109*, 8–11. [[CrossRef](#)]
57. Volovik, G.E. Comment to the CPT-symmetric Universe: Two possible extensions. *JETP Lett.* **2019**, *109*, 682–683. [[CrossRef](#)]
58. Christodoulou, M.; Riello, A.; Rovelli, C. How to detect anti-spacetime. *Int. J. Mod. Phys. D* **2012**, *21*, 1242014. [[CrossRef](#)]
59. Boyle, L.; Finn, K.; Turok, N. CPT-symmetric Universe. *Phys. Rev. Lett.* **2018**, *121*, 251301. [[CrossRef](#)] [[PubMed](#)]
60. Boyle, L.; Finn, K.; Turok, N. The Big Bang, CPT, and neutrino dark matter. *arXiv* **2018**, arXiv:1803.08930.
61. Cvetič, M.; Gibbons, G.W.; Lü, H.; Pope, C.H. Killing horizons: Negative temperatures and entropy super-additivity. *Phys. Rev. D* **2018**, *98*, 106015. [[CrossRef](#)]
62. Jacobson, T. Black hole entropy and induced gravity. *arXiv* **1994**, arXiv:gr-qc/9404039.

© 2020 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).