This is an electronic reprint of the original article.
This reprint may differ from the original in pagination and typographic detail.

Borgomeo, Edoardo; Khan, Hassaan F.; Heino, Matias; Zaveri, Esha; Kummu, Matti; Brown, Casey; Jägerskog, Anders

Impact of green water anomalies on global rainfed crop yields

Published in:
Environmental Research Letters

DOI:
[10.1088/1748-9326/abc587](https://doi.org/10.1088/1748-9326/abc587)

Published: 01/12/2020

Document Version
Publisher's PDF, also known as Version of record

Published under the following license:
CC BY

Please cite the original version:
Borgomeo, E., Khan, H. F., Heino, M., Zaveri, E., Kummu, M., Brown, C., & Jägerskog, A. (2020). Impact of green water anomalies on global rainfed crop yields. *Environmental Research Letters*, 15(12), [124030]. <https://doi.org/10.1088/1748-9326/abc587>

This material is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

LETTER • OPEN ACCESS

Impact of green water anomalies on global rainfed crop yields

To cite this article: Edoardo Borgomeo *et al* 2020 *Environ. Res. Lett.* **15** 124030

View the [article online](#) for updates and enhancements.

ENVIRONMENTAL RESEARCH
INFRASTRUCTURE AND SUSTAINABILITY™

No publication charges until 2021. Submit your research iopscience.org/eris

Editor-in-Chief

Arpad Horvath, University of California, Berkeley, USA

Environmental Research Letters

LETTER

Impact of green water anomalies on global rainfed crop yields

OPEN ACCESS

RECEIVED
21 July 2020

REVISED
5 October 2020

ACCEPTED FOR PUBLICATION
28 October 2020

PUBLISHED
3 December 2020

Original content from
this work may be used
under the terms of the
[Creative Commons
Attribution 4.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the title
of the work, journal
citation and DOI.

Edoardo Borgomeo^{1,7} , Hassaan F Khan^{2,3,7}, Matias Heino⁴ , Esha Zaveri⁵, Matti Kummu⁴ , Casey Brown⁶
and Anders Jägerskog⁵

¹ Environmental Change Institute, University of Oxford, Oxford OX1 3QY, United Kingdom

² School of Earth, Energy & Environmental Sciences, Stanford University, Stanford, CA 94305, United States of America

³ Dhanani School of Science and Engineering, Habib University, Karachi, Pakistan

⁴ Water & Development Research Group, Aalto University, P.O. Box 15200, Aalto, Finland

⁵ Water Global Practice, World Bank, 1818 H St NW, Washington, DC 20433, United States of America

⁶ Department of Civil and Environmental Engineering, University of Massachusetts Amherst, Amherst, MA 01003, United States of America

⁷ Contributed equally to the work

E-mail: edoardo.borgomeo@ouce.ox.ac.uk

Keywords: green water, hydrological variability, panel data analysis, crop yields, food production, soil moisture, rainfed agriculture

Supplementary material for this article is available [online](#)

Abstract

The importance of green water (moisture from rain stored in soils) for global food and water security is widely recognized, with process-based simulation models and field-level studies demonstrating its role in supporting rainfed agriculture. Despite this evidence, the relationship between green water anomalies and rainfed agriculture has not yet been investigated using statistical models that identify a causal relationship between the variables. Here, we address this gap and use disaggregated statistical regression (panel data analysis) at the 30 arc-min grid level to study the response of observed yields (1982–2010) of four main crops (maize, rice, soybean and wheat) to green water anomalies globally over rainfed areas. Dry green water anomalies (1 or 2 standard deviations below long-term average) decrease rainfed crop yields worldwide. This effect is more pronounced for wheat and maize, whose yields decline by 12%–18% and 7%–12% respectively. Globally, agricultural production benefits from wet green water anomalies. This effect is intensified in arid climates and weakened in humid climates where, for wheat, soybean and rice, periods of green water availability 2 standard deviations above long-term averages lead to declines in crop yield. This confirms existing evidence that excess soil moisture is detrimental to crop yield. These findings (1) advance our understanding of the impact of green water on rainfed food production and (2) provide empirical evidence supporting arguments for better management of local green water resources to reduce the impact of agricultural drought and waterlogging on rainfed crop production and capture the yield increasing effects of positive green water anomalies.

1. Introduction

Agriculture is under significant pressure. At least 800 million people are chronically undernourished, population growth and changes in consumption patterns are projected to increase global food demands, and climate change is already affecting agricultural production (FAO 2017). Against this backdrop, eradicating chronic food insecurity and malnutrition as part of the Sustainable Development Goals and meeting global food demands requires various measures to sustainably secure food supplies, including

reduced food losses, diet change and yield gap closure (Kummu *et al* 2017).

Water resources are at the center of this food security challenge and improved farm water management is an important part of closing yields gaps (Jägermeyr *et al* 2016). Increased competition for scarce surface and groundwater resources (known as ‘blue’ water) limits the potential to expand irrigation (Strzepek and Boehlert 2010) and blue water depletion could cause the conversion of agricultural production from irrigated to rainfed (Elliott *et al* 2014). Climate change, through increases in hydro-climatic

variability and extremes, will also pose challenges to agricultural production (Schmidhuber and Tubiello 2007, Porter *et al* 2014).

In this context, supporting rainfed agriculture and understanding its vulnerabilities to hydroclimatic variability are essential to ensure stability of food supplies under climate change (Cooper *et al* 2008, Rockström and Falkenmark 2015). Given the share of the world's smallholder farmers who rely on rainfed agriculture for their livelihoods, advancing our understanding of its vulnerabilities is also important to improve nutritional outcomes and livelihood opportunities for the rural poor (Rockström *et al* 2010). In addition, more stable rainfed agricultural production can reduce demand for irrigated production and allow for a reduction of irrigation water withdrawals, thus relieving pressures on scarce 'blue' water resources (Unver *et al* 2017).

Rainfed crop production depends on green water, which is here defined as moisture from rain stored in soils, following Rockström and Falkenmark (2015). Since the landmark work of Falkenmark and Rockström (Falkenmark and Rockstrom 2006), several studies have called for improved green water management—for instance through water harvesting and conservation tillage practices—to increase food security and improve rural livelihoods (Rockström *et al* 2007, Rost *et al* 2009, Sposito 2013, Rockström and Falkenmark 2015, Schyns *et al* 2019). So far, the importance of green water for crop yield has primarily been assessed using process-based simulation models, which dynamically reproduce processes affecting crop yields, and agronomic field studies.

Process-based crop simulation models have been applied to demonstrate that green water is the dominant source of water used in food production (Rost *et al* 2008, Rockström *et al* 2009a) and show that green water contributes about 90% of all water consumed by agriculture (Rost *et al* 2008, Liu *et al* 2009). Crop simulation models have also been used to assess the impacts of climate change and agricultural technologies on rainfed agriculture (Kang *et al* 2009, Calzadilla *et al* 2013, Rosegrant *et al* 2014). Siebert and Döll (2010) use a global crop water model to assess the amount of green water embedded in crop production, confirming the dominance of green water in agricultural water use worldwide. The relationship between crop yield and green water has also been studied in agronomic and field studies carried out in different parts of the world; for instance, in east and southern Africa (Rockström *et al* 2009b), China (Zhang *et al* 2004) and Spain (Lampurlanes *et al* 2016).

Despite this extensive body of work using process-based crop models, there is a limited understanding of the relationship between green water and crop yield using statistical models that are equipped to identify a causal relationship between the variables. While process-based crop models allow for attributing yield

change to specific environmental factors, they are not directly based on observational data so they might not fully reflect yield responses under real-world conditions farmers face (Ortiz-Bobea *et al* 2019). Hence, integrating findings from crop models with evidence from statistical models can provide additional insights on the impacts of hydroclimatic variables on agriculture (i.e. they are complementary methods, see Lobell and Asseng 2017, Leng and Hall 2020). A number of studies have used statistical models to show the effects of rainfall variability and other hydroclimatic variables on agricultural production (Lobell and Asseng 2017, Damania *et al* 2017, Zaveri *et al* 2020). Ortiz-Bobea *et al* (2019) and Rigden *et al* (2020) improve upon previous statistical characterizations of the yield-water relationship by highlighting the critical role of soil moisture in explaining variation in US crop yields. In this paper, we provide, to our knowledge, the first global-level estimate of the sensitivity of rainfed crop yields to green water anomalies (deviations from long-term average green water availability) using statistical models. Compared to rainfall, green water is a more precise indicator of root-zone soil moisture available for uptake by plants and is thus a better indicator of water directly available for plant growth and its impact on crop production (Falkenmark 2013).

We use disaggregated statistical regression (panel data analysis) analysis to assess how much of the observed historical interannual variation in rainfed crop yield is associated with green water anomalies. This allows us to quantify where, and how strongly, maize, rice, soybean and wheat yields respond to green water anomalies. The findings shed light on the relative importance of green water anomalies, and thus green water management, on rainfed agricultural production.

2. Data

2.1. Green water data

Following previous studies (Rockström *et al* 2009a, Kummu *et al* 2014), we use model-based estimates from the Lund-Potsdam-Jena managed Land model (LPJmL) to quantify green water. The LPJmL model was forced with the PGMFD (Princeton's Global Meteorological Forcing Dataset) global reanalysis and observation dataset (Sheffield *et al* 2006), and the simulated green water data were accessed through the Inter-Sectoral Impact Model Intercomparison Project (ISI-MIP) portal. The process-based, dynamic global vegetation and water balance model LPJmL simulates green water consumption at a daily time step on a global $0.5^\circ \times 0.5^\circ$ grid (Bondeau *et al* 2007, Rost *et al* 2008). LPJmL simulates green water consumption over both rainfed and irrigated lands. On rainfed areas, all consumed water is green water by definition so that green water consumption equals green water availability (Kummu *et al* 2014). On irrigated

areas, the available green water is defined as total evapotranspiration minus the evapotranspiration of irrigation water (i.e. blue water). Thus, to isolate the effects of green water availability on rainfed crop yield and avoid taking into account the effects of irrigation (blue water), irrigated areas are not considered in this study (see section 2.2).

To study the effects of changes in green water on rainfed crop yield, we define a green water *anomaly* using annual green water z-scores for each grid-cell, calculated over the time period (1982–2010). A dry (wet) green water anomaly in a given year occurs if green water availability in that year is at least one standard deviation lower (higher) than the long-term mean for that cell. Agricultural green water consumption refers to the water that is transpired or incorporated by the plant as well as the water that evaporates directly from the soil or leaves. Hence, a waterlogging situation can result in a positive green water anomaly due to increased availability of evaporative water. Values of -1 and 1 standard deviations are used as thresholds to define dry and wet anomalies respectively. We also examine the impact of more extreme dry and wet green water anomalies, defined using thresholds of -2 and 2 standard deviations respectively (i.e. an anomaly that is at least two standard deviations lower or higher than the long-term mean for that cell). Finally, we employ data on temperature at the grid level from Willmott and Matsuura (2001). While we do not focus on the impacts of temperature, we include it in the analysis in order to obtain unbiased estimates of the effects of green water anomalies as explained in section 3.

2.2. Rainfed crop yields

We use annual rasterized ($0.5^\circ \times 0.5^\circ$ grid) maize, rice, soybean and wheat yield (ton per hectare) data, GDHYv1.2, obtained from the Japanese Data Integration & Analysis System for years 1982–2010 (Iizumi *et al* 2014b). The crop yield data used here are based on country level agricultural statistics, which are downscaled to grid cells utilizing satellite-based net primary productivity estimates. This data set has already been used in several studies about crop yield variability (Iizumi *et al* 2014a, Iizumi and Ramankutty 2016, Iizumi *et al* 2018). Since the green water data from LPJmL (section 2.1) is mainly for a single growing season, we use crop yield data for the major growing season for maize and rice. For wheat, we use data for both spring and winter growing season, as they are mostly grown in different places, while for soybean, crop yield data is provided for a single growing season only.

To isolate crop yield data in rainfed areas, we mask the global crop yield data using information about rainfed and irrigated harvested areas from MIRCA2000 (Portmann *et al* 2010). To account for temporal changes in irrigated areas, we scale the proportion of irrigated harvested areas with information about changes (relative to year 2000) in all irrigated areas using Historical Irrigation Dataset (HID) (Siebert *et al* 2015), assuming no significant changes after year 2005, which is the last year of the data set. If the extent of annual irrigated crop specific harvested area was less than 10% of total harvested area, we consider the raster cell to be

rainfed during that year. Figure 1 shows the resulting rainfed cropland on a global $0.5^\circ \times 0.5^\circ$ grid for year 2000.

2.3. Aridity classification

We complement the crop yield and green water data with information on aridity data of each gridcell. This allows us to classify grid cells depending on their aridity index and explore the sensitivity of crop yield to green water anomalies under different aridity conditions. To differentiate grid cells based on their aridity, we use the Global Aridity Index and Potential Evapotranspiration Climate Database (Trabucco and Zomer 2019). The aridity index represents the ratio of mean annual precipitation over mean annual potential evapotranspiration for the 1970–2000 period. We define three aridity classes, as shown in figure 2.

3. Methods

We use panel data analysis (i.e. disaggregated statistical regression) to understand how green water anomalies impact rainfed crop yield of maize, rice, soybean and wheat under arid, semi-arid and humid moisture regimes. Panel data analysis is a statistical method to analyze the relationship between 2D data sets consisting of multiple observations (time series data on rainfed crop yield and green water in this case) recorded over the same sampling unit (grid cells in this case). Panel data analysis has been widely used to study the links between hydroclimatic variables and crop yields (Schlenker *et al* 2006, Dell *et al* 2014, Fishman 2016, Damania *et al* 2017, Blanc and Schlenker 2017, Zaveri and Lobell 2019, Ortiz-Bobea *et al* 2019) and also

the impact of hydroclimatic variability on economic growth (Barrios *et al* 2010, Brown *et al* 2011, Hall *et al* 2014, Sadoff *et al* 2015, Khan *et al* 2017, Gilmont *et al* 2018, Russ 2020, Damania *et al* 2020).

Our method relies on the fact that green water anomalies (the independent variable) can be considered as exogenous with respect to crop yield (the dependent variable). This holds because following standard practice in the literature, the model includes cell-fixed effects that account for unobserved, time-invariant factors that are specific to each grid cell such as soil type or other time-invariant geographic and socio-economic characteristics. It also includes time fixed effects to account for any common trends such as economic or population growth. In certain specifications (see table S2 (available online at <https://stacks.iop.org/ERL/15/124030/mmedia>)), it also includes country-specific time trends to reflect the substantial variation in technological progress across countries as well as other country-level trends. In this way, the estimation is based on random deviations of green water from its long-term mean within each grid cell, which facilitates causal inference. In order to understand the impact of green water anomalies on crop yield, the analysis estimates a log-linear model in which the outcome variable is the logarithm of yield. We estimate the following equation:

$$\log(Y)_{it} = \alpha_1 + \alpha_2 GW_{it}^- + \alpha_3 GW_{it}^+ + X'_{it}\lambda + f_c(t) + \theta_i + \gamma_i + \varepsilon_{it}. \quad (1)$$

where $\log(Y)_{it}$ is the log of rainfed crop yield in grid cell i in year t , GW_{it}^- (GW_{it}^+) is a binary variable indicating if green water was at least 1 standard deviation below (above) the long-run mean in grid cell i and

year t , $f_c(t)$ are country-specific time trends, θ_t are the year fixed effects and γ_i are the grid cell fixed effects and X'_{it} is a vector of control variables which includes log of population, and a quadratic term for mean annual temperature ($^{\circ}\text{C}$). ε_{it} represents the stochastic error term that captures variation in crop yield unexplained by the other factors in the equation. In order to obtain unbiased estimates of the effects of green water anomalies, we control for temperature. This is important since temperature has been shown to impact crop yield and the agricultural sector in general (Schlenker *et al* 2006, Schlenker and Lobell 2010, Zaveri and Lobell 2019). For each coefficient we estimate standard errors through the robust covariance matrix estimation to account for heteroskedasticity, serial correlation and cross-sectional dependence. To estimate the impact of extreme dry and wet green water anomalies, we modify this setup slightly so that GW_{it}^- (GW_{it}^+) becomes a binary variable indicating if green water was at least 2 standard deviation below (above) the long-run mean in grid cell i and year t . Summary statistics for these variables are given in table S1.

4. Results and interpretation

We examine where and by how much green water anomalies impact rainfed crop yields. At the global level, we find that rainfed crop yields increase in response to wet green water anomalies and decrease with dry green water anomalies across the four crops considered, as shown in figure 3. These effects are statistically significant for both wet and dry anomalies and are robust to alternate specifications, meaning that the significance of the impact and magnitude of the coefficient remains largely unchanged in the specifications (see table S2). Wheat and maize show the stronger sensitivity to both wet and dry anomalies compared to rice and soybean. This provides empirical evidence to support the arguments that green water deficit (i.e. agricultural drought) damages rainfed agricultural production and food security globally, and that additional soil moisture enhances crop yields (Rockström and Falkenmark 2015).

The effects of green water anomalies vary under aridity conditions. For all four crops, the negative effect of dry anomalies is weakened for the humid areas and intensified for the semi-arid and arid areas, as shown in figure 4 and table S3. Similarly, the effect of wet anomalies is greater for the arid and semi-arid areas. This suggests that additional green water is beneficial to crop yields especially in arid regions, which also explains the widespread use of supplementary irrigation in these areas. For the humid sample, we find a null or slightly positive effect of 1 standard deviation wet anomalies on crop yields except for wheat that shows negative effect (figure 4). When we examine extreme wet anomalies in humid regions (2 standard deviations above long-run mean),

we find a negative effect on crop yields for wheat, soybean and rice. At first, this result may seem counterintuitive as increased soil moisture availability, and thus water uptake by plants, is anticipated to increase crop yield. However, excess soil moisture can have negative direct and indirect effects on crop growth in humid climates. Excess soil moisture (so a wet green water anomaly) can be directly damaging to crops because it creates anoxic conditions in soils, and it can also have indirect effects, increasing the risk of plant diseases and pests and delaying farmers' planting and harvesting activities (Rosenzweig *et al* 2002). These results are aligned with crop-specific findings in the literature which we discuss next.

4.1. Maize

We find that dry green water anomalies have a statistically significant negative (decreasing) effect on rainfed maize yields globally (figure 3). When we disaggregate the results by aridity class (figure 4), we find that the negative (decreasing) effects of dry green water anomalies on maize yield are particularly pronounced in arid and semi-arid regions (23%–39% reduction in yields). Field studies from several arid countries confirm our finding that rainfed maize yield in arid areas is highly sensitive to below-average soil moisture levels (e.g. Rockström *et al* 2009b). In semi-arid parts of China, simulation studies demonstrate the impact of water shortage on rainfed maize, highlight the increasing importance of water as a limiting factor partly because of the shift to longer-maturing maize varieties and agronomic practices (e.g. plant density which increase water demand) (Meng *et al* 2016). Finally, our finding that dry green water anomalies negatively impact maize yields in humid climates is in agreement with statistical (Rigden *et al* 2020) and modelling (Andresen *et al* 2001) analyses from Midwest USA and also, Germany (statistical analysis from Peichl *et al* 2018), which both fall within our humid climate category (see figure 2).

4.2. Wheat

When green water is 1 standard deviation below the long-run mean, rainfed wheat yield decreases globally by 12%–15% (figure 3). This effect is larger in arid and semi-arid areas (figure 4), in agreement with Daryanto *et al* (2016)'s global synthesis, where wheat sensitivity to drought in arid areas is found to be greater than in humid areas. Given the low relative humidity and high potential evapotranspiration in arid areas, this finding is not surprising. As shown in figure 4, wheat also shows the lowest sensitivity to dry green water anomalies in humid areas across the four crops considered. This could be in part attributed to the differences in origin and traits between wheat and the other crops, such as maize. While maize is thought to originate from humid areas (Van Heerwaarden *et al* 2011), wheat originates from dry regions (Charmet

2011), and could thus show lower sensitivity to drought in humid regions (Daryanto *et al* 2016).

Globally, additional green water has a positive effect on rainfed wheat yields, with a 1 standard deviation wet anomaly increasing yield by 5% (figure 3). This is line with evidence suggesting that higher yields are obtained when wheat is irrigated (e.g. more than double under certain conditions, Cao *et al* 2014) or grown under soil moisture conservation practices (e.g. mulching in no-till fields, Rahman *et al* 2005, Guo *et al* 2019). However, the positive effect of excess green water is reversed in humid countries especially

for extreme wet anomalies (2 standard deviations above the long-term mean) which reduce yield by 7% (figure 4). In humid regions, this effect is larger than the effect of droughts, suggesting that excess soil moisture is a key factor explaining rainfed wheat yield changes in areas such as the eastern USA and parts of South America. This latter effect is in agreement with other assessments of wheat sensitivity to hydro-climatic shocks; for instance, Zampieri *et al* (2017) using a statistical approach based on historical data found that excess water availability negatively affects wheat production in humid areas of the world.

4.3. Soybean

At the global level, soybean is sensitive to green water anomalies (figure 3). This confirms evidence from agronomic studies (Foroud *et al* 1993, Wijewardana *et al* 2018) and also crop models (e.g. Bhatia *et al* 2008) showing that soil moisture is a major factor limiting yield of rainfed soybean. Globally, a dry green water anomaly leads to soybean yield reductions between 6% to 12%, depending on the size of the anomaly. This confirms other statistical assessments of the effects of drought on observed soybean yields (e.g. Troy *et al* 2015, Leng *et al* 2016, Zipper *et al* 2016).

Additional soil moisture boosts rainfed soybean yields, as also suggested by field studies (Dass and Bhattacharyya 2017). However, in humid areas, when excess soil moisture is 2 standard deviations higher than the long-term average, yields show a modest (−1%) decrease. This confirms soybean sensitivity to water-logging, and can be explained by considering the direct and indirect effects of excess soil moisture on crop growth and farmers' decisions (e.g. delaying farmers' planting and harvesting activities) (Rosenzweig *et al* 2002, Rhine *et al* 2010).

4.4. Rice

Globally, rice shows the lowest sensitivity to green water anomalies among the four crops considered (figure 3). For both dry and wet anomalies in soil moisture, the magnitude of the effect on rainfed rice yield is similar, at about 2%. However, the magnitude of this effect increases sizably when we disaggregate the results by aridity (figure 4). In arid and semi-arid areas, agricultural drought reduces rice yields by 5%–23%, under a 1 and 2 standard deviation anomaly, respectively. This is not surprising because under rainfed conditions in arid and semi-arid climates (e.g. some parts of Nigeria and China for example) rice fields are intermittently flooded depending on water availability (Steduto *et al* 2012), making rainfed rice yield particularly sensitive to water deficit (Lilley and Fukai 1994, Pandey *et al* 2000). We also find that low soil moisture leads to declines in rainfed rice yields in humid areas, in agreement with findings from humid countries (e.g. observed data from Philippines in Stuecker *et al* 2018).

Unlike the other cereals considered in this study, rice grows well in paddy fields and is known to be tolerant to excess water. We find that moderate excess water (positive 1 standard deviation anomaly) has a positive effect on rainfed rice yields under all aridity conditions (figure 4). This positive effect is also observed for the 2 standard deviation positive anomaly, apart for humid areas, where it becomes negative. This suggests that, despite having tolerance for excess water, rainfed rice is still sensitive to excess green water in humid areas (i.e. severe waterlogging associated with floods). Potential mechanisms to explain this result include elongation growth during flash flooding which competes with maintenance processes requiring carbohydrates and energy, thus negatively affecting plant growth and survival (Ram *et al* 2002).

5. Discussion and conclusions

This study presents the first global statistical analysis of the effects of green water anomalies on observed crop yields for four major crops. The results show that dry green water anomalies (soil moisture deficit) have a statistically significant negative (decreasing) effect on rainfed crop yields globally. This effect is larger than that of wet green water anomalies, which globally have a positive (increasing) effect on yields. The analysis demonstrates the greater sensitivity of certain crops to green water variability, with wheat and maize showing the greatest sensitivity to dry green water anomalies and wet green water anomalies compared to soybean and rice. Excess soil moisture has a negative impact on rainfed wheat and soybean crop yields, highlighting the importance for research to focus not just on green water impacts under droughts but also under extreme wet conditions.

As all statistical assessments of the effects of hydroclimatic variables on crop production (e.g. Vogel *et al* 2019), our results hinge upon our data and methodological choices. The temporal resolution for the crop yield and green water data is growing season, which may be too low to model specific green water and crop growth relationships. This means that we are unable to understand if green water anomalies are only important during particular phases of the plant's growing season. We did not explore the sensitivity of crop yields to green water anomalies for specific stages of crop growth and for different crop growth processes, which are known to be important factors in determining crop yield responses (Siebert *et al* 2017a). To address these limitations, future research should seek to complement the empirical results presented in this study with process-based crop model simulations to differentiate among drivers (e.g. Heino *et al* 2018) and also test the effects of improved green water management practices and adoption of technologies on reducing crop yield fluctuations (e.g. Frieler *et al* 2017, Siebert *et al* 2017b). Advancing this understanding is a key to support efforts aimed at improving food security and incomes of the millions of people living in rainfed areas.

Our findings provide empirical evidence to support arguments that green water management is important for global food production and can enhance agricultural resilience to dry spells (Rockström and Falkenmark 2015). Measures to retain more water in soils (e.g. use of cover crops, no-till systems, soil water management practices such as terraces), thus artificially augmenting green water, would contribute to making rainfed crop production more stable in the face of green water deficit and agricultural drought (Basche *et al* 2016, Lampurlanes *et al* 2016, Kosmowski 2018). Despite the positive effects of these measures on yields (Rockström and Falkenmark 2015), their benefits often depend on a host of other factors and other measures such as crop rotation, residue retention, and aridity (Pittelkow *et al* 2015), which have to be considered when promoting green water management practices.

Acknowledgments

We acknowledge ISI-MIP for providing and sharing the green water data used in this study. MK received financial support from Strategic Research Council (SRC) through project 'From Failand to Winland', Emil Aaltonen foundation through project 'eat-less-water', European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant agreement No. 819202), and Academy of Finland funded project WATVUL (Grant No. 317320). MH was financially

supported by Maa- ja vesitekniiikan tuki ry, the Vilho, Yrjö and Kalle Väisälä Foundation and AaltoENG doctoral programme. We thank three reviewers for their comments which helped to greatly improve the paper. The processing scripts are available from: <https://github.com/matheino/greenwater>.

Data availability statement

The data that support the findings of this study are openly available at the following URL/DOI: <https://github.com/matheino/greenwater>.

ORCID iDs

Edoardo Borgomeo <https://orcid.org/0000-0002-8351-9064>

Matias Heino <https://orcid.org/0000-0003-4294-7756>

Matti Kummu <https://orcid.org/0000-0001-5096-0163>

References

- Andresen J A, Alagarswamy G, Rotz C A, Ritchie J T and Lebaron A W 2001 Weather impacts on maize, soybean, and alfalfa production in the Great Lakes region, 1895–1996 *Agron. J.* **93** 1059–70
- Barrios S, Bertinelli L and Strobl E 2010 Trends in rainfall and economic growth in Africa: a neglected cause of the African growth tragedy *Rev. Econ. Stat.* **92** 350–66
- Basche A D, Kaspar T C, Archontoulis S V, Jaynes D B, Sauer T J, Parkin T B and Miguez F E 2016 Soil water improvements with the long-term use of a winter rye cover crop *Agric. Water Manag.* **172** 40–50
- Bhatia V S, Singh P, Wani S P, Chauhan G S, Rao A K, Mishra A K and Srinivas K 2008 Analysis of potential yields and yield gaps of rainfed soybean in India using CROPGRO-Soybean model *Agric. For. Meteorol.* **148** 1252–1265
- Blanc E and Schlenker W 2017 The use of panel models in assessments of climate impacts on agriculture *Rev. Environ. Econ. Policy* **11** 258–79
- Bondeau A et al 2007 Modelling the role of agriculture for the 20th century global terrestrial carbon balance *Glob. Chang. Biol.* **13** 679–706
- Brown C, Meeks R, Hunu K and Yu W 2011 Hydroclimate risk to economic growth in sub-Saharan Africa *Clim. Change* **106** 621–47
- Calzadilla A, Rehdanz K, Betts R, Falloon P, Wiltshire A and Tol R S 2013 Climate change impacts on global agriculture *Clim. Change* **120** 357–74
- Cao X C, Wu P T, Wang Y B and Zhao X N 2014 Assessing blue and green water utilisation in wheat production of China from the perspectives of water footprint and total water use *Hydrol. Earth Syst. Sci.* **18** 3165
- Charmet G 2011 Wheat domestication: lessons for the future *C. R. Biol.* **334** 212–20
- Cooper P J M, Dimes J, Rao K P C, Shapiro B, Shiferaw B and Twomlow S 2008 Coping better with current climatic variability in the rain-fed farming systems of sub-Saharan Africa: an essential first step in adapting to future climate change? *Agric. Ecosyst. Environ.* **126** 24–35
- Damania R et al 2017 *Uncharted Waters: The New Economics of Water Scarcity and Variability* (Washington, DC: World Bank)
- Damania R, Desbureaux S and Zaveri E 2020 Does rainfall matter for economic growth? Evidence from global sub-national data (1990–2014) *J. Environ. Econ. Manag.* **102** 102335
- Daryanto S, Wang L and Jacinthe P A 2016 Global synthesis of drought effects on maize and wheat production *PloS One* **11** e0156362
- Dass A and Bhattacharyya R 2017 Wheat residue mulch and anti-transpirants improve productivity and quality of rainfed soybean in semi-arid north-Indian plains *Field Crops Res.* **210** 9–19
- Dell M, Jones B F and Olken B A 2014 What do we learn from the weather? The new climate-economy literature *J. Econ. Lit.* **52** 740–98
- Elliott J et al 2014 Constraints and potentials of future irrigation water availability on agricultural production under climate change *Proc. Natl Acad. Sci.* **111** 3239–44
- Falkenmark M 2013 Growing water scarcity in agriculture: future challenge to global water security *Phil. Trans. R. Soc. A* **371** 20120410
- Falkenmark M and Rockstrom J 2006 The new blue and green water paradigm: breaking new ground for water resources planning and management *J. Water Resour. Plan. Manag.* **132** 129–132
- FAO 2017 *The Future of Food and Agriculture. Trends and Challenges* (Rome: FAO)
- Fishman R 2016 More uneven distributions overturn benefits of higher precipitation for crop yields *Environ. Res. Lett.* **11** 024004
- Foroud N, Mündel H H, Saindon G and Entz T 1993 Effect of level and timing of moisture stress on soybean plant development and yield components *Irrig. Sci.* **13** 149–155
- Frieler K et al 2017 Understanding the weather signal in national crop-yield variability *Earths Future* **5** 605–16
- Gilmont M, Hall J W, Grey D, Dadson S J, Abele S and Simpson M 2018 Analysis of the relationship between rainfall and economic growth in Indian states *Glob. Environ. Change* **49** 56–72
- Guo Y, Yin W, Hu F, Fan Z, Fan H, Zhao C, Yu A, Chai Q and Coulter J A 2019 Reduced irrigation and nitrogen coupled with no-tillage and plastic mulching increase wheat yield in maize-wheat rotation in an arid region *Field Crops Res.* **243** 107615
- Hall J W, Grey D, Garrick D, Fung E, Brown C, Dadson S J and Sadoff C W 2014 Coping with the curse of freshwater variability *Science* **346** 429–430
- Heino M, Puma M J, Ward P J, Gerten D, Heck V, Siebert S and Kummu M 2018 Two-thirds of global cropland area impacted by climate oscillations *Nat. Commun.* **9** 1257
- Iizumi T, Kotoku M, Kim W, West P C, Gerber J S and Brown M E 2018 Uncertainties of potentials and recent changes in global yields of major crops resulting from census-and satellite-based yield datasets at multiple resolutions *PloS One* **13** e0203809
- Iizumi T, Luo -J-J, Challinor A J, Sakurai G, Yokozawa M, Sakuma H, Brown M E and Yamagata T 2014a Impacts of El Niño Southern Oscillation on the global yields of major crops *Nat. Commun.* **5** 3712
- Iizumi T and Ramankutty N 2016 Changes in yield variability of major crops for 1981–2010 explained by climate change *Environ. Res. Lett.* **11** 034003
- Iizumi T, Yokozawa M, Sakurai G, Travasso M I, Romanenkov V, Oettli P, Newby T, Ishigooka Y and Furuya J 2014b Historical changes in global yields: major cereal and legume crops from 1982 to 2006 *Glob. Ecol. Biogeogr.* **23** 346–357
- Jägermeyr J, Gerten D, Schaphoff S, Heinke J, Lucht W and Rockström J 2016 Integrated crop water management might sustainably halve the global food gap *Environ. Res. Lett.* **11** 025002
- Kang Y, Khan S and Ma X 2009 Climate change impacts on crop yield, crop water productivity and food security—a review *Prog. Nat. Sci.* **19** 1665–74
- Khan H F, Morzuch B J and Brown C M 2017 Water and growth: an econometric analysis of climate and policy impacts *Water Resour. Res.* **53** 5124–36
- Kosmowski F 2018 Soil water management practices (terraces) helped to mitigate the 2015 drought in Ethiopia *Agric. Water Manag.* **204** 11–16

- Kummu M, Fader M, Gerten D, Guillaume J H, Jalava M, Jägermeyr J, Pfister S, Porkka M, Siebert S and Varis O 2017 Bringing it all together: linking measures to secure nations' food supply *Curr. Opin. Environ. Sustain.* **29** 98–117
- Kummu M, Gerten D, Heinke J, Konzmann M and Varis O 2014 Climate-driven interannual variability of water scarcity in food production potential: a global analysis *Hydrol. Earth Syst. Sci.* **18** 447–61
- Lampurlanes J, Plaza-Bonilla D, Alvaro-Fuentes J and Cantero-Martinez C 2016 Long-term analysis of soil water conservation and crop yield under different tillage systems in Mediterranean rainfed conditions *Field Crops Res.* **189** 58–67
- Leng G and Hall J W 2020 Predicting spatial and temporal variability in crop yields: an inter-comparison of machine learning, regression and process-based models *Environ. Res. Lett.* **15** 044027
- Leng G, Zhang X, Huang M, Asrar G R and Leung L R 2016 The role of climate covariability on crop yields in the conterminous United States *Sci. Rep.* **6** 33160
- Lilley J M and Fukai S 1994 Effect of timing and severity of water deficit on four diverse rice cultivars II. Physiological responses to soil water deficit *Field Crops Res.* **37** 215–23
- Liu J, Zehnder A J B and Yang H 2009 Global consumptive water use for crop production: the importance of green water and virtual water *Water Resour. Res.* **45** W05428
- Lobell D B and Asseng S 2017 Comparing estimates of climate change impacts from process-based and statistical crop models *Environ. Res. Lett.* **12** 015001
- Meng Q, Chen X, Lobell D B, Cui Z, Zhang Y, Yang H and Zhang F 2016 Growing sensitivity of maize to water scarcity under climate change *Sci. Rep.* **6** 19605
- Ortiz-Bobea A, Wang H, Carrillo C M and Ault T R 2019 Unpacking the climatic drivers of US agricultural yields *Environ. Res. Lett.* **14** 064003
- Pandey S, Behura D D, Villano R and Naik D 2000 Economic cost of drought and farmers' coping mechanisms: a study of rainfed rice systems in eastern India International Rice Research Institute *IRRI Discussion Papers* 287599
- Peichl M, Thober S, Meyer V and Samaniego L 2018 The effect of soil moisture anomalies on maize yield in Germany *Nat. Hazards Earth Syst. Sci.* **18** 889–906
- Pittelkow C M et al 2015 Productivity limits and potentials of the principles of conservation agriculture *Nature* **517** 365–8
- Porter J, Xie L, Challinor A, Cochrane K, Howden S, Iqbal M, Lobell D and Travasso M 2014 Food security and food production systems *Climate Change 2014: Impacts, Adaptation, and Vulnerability: A. Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed C Field et al (Cambridge: Cambridge University Press) pp 485–533
- Portmann F T, Siebert S and Döll P 2010 MIRCA2000—Global monthly irrigated and rainfed crop areas around the year 2000: a new high-resolution data set for agricultural and hydrological modeling *Glob. Biogeochem. Cycles* **24** 1–24
- Rahman M A, Chikushi J, Saifuzzaman M and Lauren J G 2005 Rice straw mulching and nitrogen response of no-till wheat following rice in Bangladesh *Field Crops Res.* **91** 71–81
- Ram P C et al 2002 Submergence tolerance in rainfed lowland rice: physiological basis and prospects for cultivar improvement through marker-aided breeding *Field Crops Res.* **76** 131–52
- Rhine M D, Stevens G, Shannon G, Wrather A and Sleper D 2010 Yield and nutritional responses to waterlogging of soybean cultivars *Irrig. Sci.* **28** 135–42
- Rigden A J, Mueller N D, Holbrook N M, Pillai N and Huybers P 2020 Combined influence of soil moisture and atmospheric evaporative demand is important for accurately predicting US maize yields *Nat. Food* **1** 127–33
- Rockström J et al 2009b Conservation farming strategies in East and Southern Africa: yields and rain water productivity from on-farm action research *Soil Tillage Res.* **103** 23–32
- Rockström J and Falkenmark M 2015 Increase water harvesting in Africa *Nature* **519** 283–5
- Rockström J, Falkenmark M, Karlberg L, Hoff H, Rost S and Gerten D 2009a Future water availability for global food production: the potential of green water for increasing resilience to global change *Water Resour. Res.* **45** W00A12
- Rockström J, Karlberg L, Wani S P, Barron J, Hatibu N, Oweis T, Bruggeman A, Farahani J and Qiang Z 2010 Managing water in rainfed agriculture—the need for a paradigm shift *Agric. Water Manag.* **97** 543–550
- Rockström J, Lannerstad M and Falkenmark M 2007 Assessing the water challenge of a new green revolution in developing countries *Proc. Natl Acad. Sci.* **104** 6253–60
- Rosegrant M W et al 2014 Food security in a world of natural resource scarcity: the role of agricultural technologies *Intl. Food Policy Res. Inst.*
- Rosenzweig C, Tubiello F N, Goldberg R, Mills E and Bloomfield J 2002 Increased crop damage in the US from excess precipitation under climate change *Glob. Environ. Change* **12** 197–202
- Rost S, Gerten D, Bondeau A, Lucht W, Rohwer J and Schaphoff S 2008 Agricultural green and blue water consumption and its influence on the global water system *Water Resour. Res.* **44** W09405
- Rost S, Gerten D, Hoff H, Lucht W, Falkenmark M and Rockström J 2009 Global potential to increase crop production through water management in rainfed agriculture *Environ. Res. Lett.* **4** 044002
- Russ J 2020 Water runoff and economic activity: the impact of water supply shocks on growth *J. Environ. Econ. Manag.* **101** 102322
- Sadoff C W et al 2015 *Securing Water, Sustaining Growth: Report of the GWP/OECD Task Force on Water Security and Sustainable Growth* University of Oxford, Oxford p 180
- Schlenker W, Hanemann W M and Fisher A C 2006 The impact of global warming on US agriculture: an econometric analysis of optimal growing conditions *Rev. Econ. Stat.* **88** 113–25
- Schlenker W and Lobell D B 2010 Robust negative impacts of climate change on African agriculture *Environ. Res. Lett.* **5** 014010
- Schmidhuber J and Tubiello F N 2007 Global food security under climate change *Proc. Natl Acad. Sci.* **104** 19703–8
- Schyns J F, Hoekstra A Y, Booij M J, Hogeboom R J and Mekonnen M M 2019 Limits to the world's green water resources for food, feed, fiber, timber, and bioenergy *Proc. Natl Acad. Sci.* **116** 4893–8
- Sheffield J, Goteti G and Wood E F 2006 Development of a 50-year high-resolution global dataset of meteorological forcings for land surface modeling *J. Clim.* **19** 3088–111
- Siebert S and Döll P 2010 Quantifying blue and green virtual water contents in global crop production as well as potential production losses without irrigation *J. Hydrol.* **384** 198–217
- Siebert S, Kummu M, Porkka M, Döll P, Ramankutty N and Scanlon B R 2015 A global data set of the extent of irrigated land from 1900 to 2005 *Hydrol. Earth Syst. Sci.* **19** 1521–45
- Siebert S, Webber H and Rezaei E E 2017a Weather impacts on crop yields—searching for simple answers to a complex problem *Environ. Res. Lett.* **12** 081001
- Siebert S, Webber H, Zhao G and Ewert F 2017b Heat stress is overestimated in climate impact studies for irrigated agriculture *Environ. Res. Lett.* **12** 054023
- Sposito G 2013 Green water and global food security *Vadose Zone J.* **12** 1–6
- Steduto P, Hsiao T C, Fereres E and Raes D 2012 *Crop Yield Response to Water* vol 1028 (Rome: FAO)
- Strzepek K and Boehlert B 2010 Competition for water for the food system *Phil. Trans. R. Soc. B* **365** 2927–40
- Stuecker M F, Tigchelaar M and Kantar M B 2018 Climate variability impacts on rice production in the Philippines *PloS One* **13** e0201426

- Trabucco A and Zomer R 2019 Global aridity index and potential evapotranspiration (ET0) climate database v2. figshare *Fileset* (available at: <https://doi.10.6084/m9.figshare.7504448.v3>)
- Troy T J, Kipgen C and Pal I 2015 The impact of climate extremes and irrigation on US crop yields *Environ. Res. Lett.* **10** 054013
- Unver O, Bhaduri A and Hoogeveen J 2017 Water-use efficiency and productivity improvements towards a sustainable pathway for meeting future water demand *Water Secur.* **1** 21–27
- Van Heerwaarden J, Doebley J, Briggs W H, Glaubitz J C, Goodman M M, Gonzalez J D J S and Ross-Ibarra J 2011 Genetic signals of origin, spread, and introgression in a large sample of maize landraces *Proc. Natl Acad. Sci.* **108** 1088–92
- Vogel E, Donat M G, Alexander L V, Meinshausen M, Ray D K, Karoly D, Meinshausen N and Frieler K 2019 The effects of climate extremes on global agricultural yields *Environ. Res. Lett.* **14** 054010
- Wijewardana C, Reddy K R, Alsajri F A, Irby J T, Krutz J and Golden B 2018 Quantifying soil moisture deficit effects on soybean yield and yield component distribution patterns *Irrig. Sci.* **36** 241–255
- Willmott C J and Matsuura K 2001 Terrestrial air temperature and precipitation: monthly and annual time series (1950–1999) (available at: http://climate.geog.udel.edu/~climate/html_pages/README.ghcn_ts2.html)
- Zampieri M, Ceglar A, Dentener F and Toreti A 2017 Wheat yield loss attributable to heat waves, drought and water excess at the global, national and subnational scales *Environ. Res. Lett.* **12** 064008
- Zaveri E and Lobell D B 2019 The role of irrigation in changing wheat yields and heat sensitivity in India *Nat. Commun.* **10** 1–7
- Zaveri E, Russ J and Damania R 2020 Rainfall anomalies are a significant driver of cropland expansion *Proc. Natl Acad. Sci.* **117** 10225–10233
- Zhang Y, Kendy E, Qiang Y, Changming L, Yanjun S and Hongyong S 2004 Effect of soil water deficit on evapotranspiration, crop yield, and water use efficiency in the North China Plain *Agric. Water Manag.* **64** 107–22
- Zipper S C, Qiu J and Kucharik C J 2016 Drought effects on US maize and soybean production: spatiotemporal patterns and historical changes *Environ. Res. Lett.* **11** 094021