This is an electronic reprint of the original article.
This reprint may differ from the original in pagination and typographic detail.

Suomela, S.; Sampaio, R.; Ala-Nissilä, Tapio

Comparison between quantum jumps and master equation in the presence of a finite environment

Published in:
Physical Review E

DOI:
[10.1103/PhysRevE.94.032138](https://doi.org/10.1103/PhysRevE.94.032138)

Published: 30/09/2016

Document Version
Publisher's PDF, also known as Version of record

Please cite the original version:
Suomela, S., Sampaio, R., & Ala-Nissilä, T. (2016). Comparison between quantum jumps and master equation in the presence of a finite environment. *Physical Review E*, 94(3), 1-9. [032138].
<https://doi.org/10.1103/PhysRevE.94.032138>

Comparison between quantum jumps and master equation in the presence of a finite environment

S. Suomela,¹ R. Sampaio,¹ and T. Ala-Nissila^{1,2}¹*Department of Applied Physics and COMP Centre of Excellence, Aalto University School of Science, P.O. Box 11100, 00076 Aalto, Finland*²*Department of Physics, P.O. Box 1843, Brown University, Providence, Rhode Island 02912-1843, USA*

(Received 22 July 2016; published 30 September 2016)

We study the equivalence between the recently proposed finite environment quantum jump model and a master equation approach. We derive microscopically the master equation for a qubit coupled to a finite bosonic environment and show that the master equation is equivalent to the finite environment quantum jump model. We analytically show that both the methods produce the same moments of work when the work is defined through the two-measurement protocol excluding the interaction energy. However, when compared to the work moments computed using the power operator approach, we find a difference in the form of the work moments. To numerically verify our results, we study a qubit coupled to an environment consisting of ten two-level systems.

DOI: [10.1103/PhysRevE.94.032138](https://doi.org/10.1103/PhysRevE.94.032138)

I. INTRODUCTION

Although the thermodynamics of small quantum systems has been intensively studied in the past few years [1–34], measuring thermodynamic variables, such as work, heat, and entropy, and their fluctuations in these systems has turned out to be a difficult task. Only recently have the work fluctuation relations been experimentally verified in driven closed quantum systems using the two-measurement protocol [13,14]. For open quantum systems the situation is even more problematic. Instead of measuring only the internal energy of the system at the beginning and at the end of the drive, also the heat emitted to the environment must be measured. One of the few proposed measurement schemes to address this issue is the calorimetric detection of the immediate environment [35].

In the calorimetric measurement, the small quantum system is coupled to a large but finite environment. The coupling between the system and the environment is assumed to be weak enough such that it can be neglected in the energy terms and modeled by stochastic jumps alone. Due to these environment-induced jumps between the system eigenstates, heat is exchanged with the environment. The resulting changes in the environment energy are continuously monitored by a detector. As a consequence, the heat released to the environment can be measured without directly measuring the quantum system. For a two-level quantum system (qubit), the internal energy change of the system can also be obtained from the heat emitted (see, for example, Appendix C in Ref. [36]). As the environment is large and coupled to a detector, it is assumed to be decohered into a set of energy eigenstates, called microstates [36].

In a recent paper [36], a finite-environment quantum jump (FEQJ) model was introduced to describe the calorimetric process. In the model, a jump changes both the system and the environment states. The evolution of the system is non-Markovian as its previous history affects its future evolution through the evolution of the environment.

In this article, we study the compatibility of the FEQJ model with the corresponding master equation approach. We focus on a weakly driven qubit coupled to a finite bosonic environment, called a calorimeter from here on. The setup is illustrated in Fig. 1. The qubit and calorimeter are assumed to be initially thermalized by an ideal bath. During the driving protocol, we

neglect the coupling to the ideal bath as it is assumed to be very weak compared to the inverse of the total driving time τ . Due to the detector, the calorimeter is assumed to decohere into its eigenstates.

We start the article by microscopically deriving the master equation for the qubit coupled to the calorimeter. We then show that the master equation obtained is equivalent to the master equation formed from the FEQJ model. We also show that both methods produce the same work moments when the work is defined using the two-measurement protocol (TMP) [37–39] without the interaction energy. For master equation calculations, we additionally show that the power operator definition of work [23,24] does not produce exactly the same work moments due to a different order of approximations. Finally, we numerically study a qubit coupled to an calorimeter consisting of ten two-level systems with an energy gap equivalent to that of the qubit. We calculate the qubit density matrix and the first two moments of work with the FEQJ method and directly from the master equation. We find excellent agreement between the methods when the work is defined using the TMP without the interaction energy. However, when compared to the work moments from the power operator approach, we show that the agreement sensitively depends on the form of the driving.

II. MICROSCOPIC DERIVATION OF THE MASTER EQUATION

We focus on a two-level system (qubit) $H_0 = \hbar\omega_0 a^\dagger a$ that is weakly driven by a classical source $V_D(t) = \lambda(t)a^\dagger + \lambda^*(t)a$, where $a = |0\rangle\langle 1|$ and $a^\dagger = |1\rangle\langle 0|$ are, respectively, the annihilation and creation operators in the undriven basis. The states $|1\rangle$ and $|0\rangle$ denote the excited and ground states of the undriven Hamiltonian H_0 , respectively. The qubit Hamiltonian is then given by $H_q(t) = H_0 + V_D(t)$. As shown in Fig. 1, the qubit is weakly coupled to a bosonic calorimeter by $V = \sum_k \kappa_k (a^\dagger d_k + a d_k^\dagger)$, where the coupling strength κ_k is real and d_k and $d_k^\dagger$ are the calorimeter's annihilation and creation operators associated with energy ϵ_k , respectively. The calorimeter Hamiltonian is given by $H_c = \sum_k \epsilon_k d_k^\dagger d_k$.

Let us start from the total density matrix of the qubit-calorimeter composite, which can always be expressed

FIG. 1. Schematic illustration of the calorimetric setup. The qubit and the calorimeter start from thermal equilibrium with an ideal bath. During the protocol, the qubit is driven by a classical field $\lambda(t)$. The calorimeter is constantly monitored with a detector. In the simulations, the calorimeter is assumed to contain two-level systems with an energy gap equivalent to that of the qubit $\hbar\omega_0$.

as

$$\rho(t) = \sum_{i,j,m,n} \alpha_{ijmn}(t) |i\rangle\langle j| \otimes |\Psi_m\rangle\langle\Psi_n|, \quad (1)$$

where $|\Psi_n\rangle$ are calorimeter energy eigenstates (microstates) forming a complete basis. In order to witness the energy changes in the calorimeter, it is continuously monitored with a detector. As a consequence of the monitoring, we assume that the calorimeter decoheres instantaneously into the environment-induced superselection basis with $\alpha_{ijmn}(t) = 0$ if $m \neq n$ [40]. Consequently, the total density matrix simplifies to

$$\rho(t) = \sum_n \sigma(n,t) \otimes |\Psi_n\rangle\langle\Psi_n|, \quad (2)$$

where $\sigma(n,t) = \sum_{i,j} \alpha_{ijnn}(t) |i\rangle\langle j|$. It should be noted that the matrix $\sigma(n,t)$ cannot be interpreted as a qubit density matrix since its trace over qubit degrees of freedom gives the probability of a calorimeter state $|\Psi_n\rangle$, denoted by $p(\Psi_n, t)$. The reduced density matrices of the qubit and the calorimeter are given by $\rho_q(t) = \text{Tr}_c\{\rho(t)\} = \sum_n \sigma(n,t)$ and $\rho_c(t) = \text{Tr}_q\{\rho(t)\}$, respectively, where the subscript c (q) in the trace denotes the trace over the calorimeter (qubit) degrees of freedom.

Let us assume that the total density matrix starts from a tensor product state of the form of Eq. (2). In the interaction picture with respect to $H_0 + H_c$, the time evolution of the total density matrix is given by the following equation up to the second order of V_D and V :

$$\begin{aligned} \dot{\rho}_I(t) &= \frac{i}{\hbar} [\rho_I(t), V_{D,I}(t)] \\ &\quad - \frac{1}{\hbar^2} \int_{-\infty}^t dt' [V_I(t), [V_{D,I}(t') + V_I(t'), \rho_I(t)']], \end{aligned} \quad (3)$$

where the subscript I denotes the interaction picture, e.g., $V_I(t) = e^{i(H_0+H_c)t/\hbar} V(t) e^{-i(H_0+H_c)t/\hbar}$. The master equation for $\sigma_I(n,t) = \text{Tr}_c\{\rho_I(t) |\Psi_n\rangle\langle\Psi_n|\}$ then takes the form

$$\begin{aligned} \dot{\sigma}_I(n,t) &= \frac{i}{\hbar} [\sigma_I(n,t), V_{D,I}(t)] \\ &\quad - \frac{1}{\hbar^2} \int_{-\infty}^t dt' \text{Tr}_c\{[V_I(t), [V_I(t'), \rho_I(t)]] |\Psi_n\rangle\langle\Psi_n|\}. \end{aligned} \quad (4)$$

By inserting the Hamiltonians and neglecting the Lamb shift (see Appendix A for details), we obtain the following master equation in the Schrödinger picture:

$$\begin{aligned} \dot{\sigma}(n,t) &= \frac{i}{\hbar} [\sigma(n,t), H_q(t)] \\ &\quad - \sum_k \left\{ \frac{\Gamma_{\uparrow,k}(n)}{2} [\sigma(n,t) a a^\dagger + a a^\dagger \sigma(n,t)] \right. \\ &\quad - \Gamma_{\downarrow,k}(n'_k) a \sigma(n'_k, t) a^\dagger - \Gamma_{\uparrow,k}(n''_k) a^\dagger \sigma(n''_k, t) a \\ &\quad \left. + \frac{\Gamma_{\downarrow,k}(n)}{2} [\sigma(n,t) a^\dagger a + a^\dagger a \sigma(n,t)] \right\}, \end{aligned} \quad (5)$$

where n'_k and n''_k are the calorimeter microstate indices that satisfy $|\Psi_{n'_k}\rangle = d_k |\Psi_n\rangle / \|d_k |\Psi_n\rangle\|$ and $|\Psi_{n''_k}\rangle = d_k^\dagger |\Psi_n\rangle / \|d_k^\dagger |\Psi_n\rangle\|$, respectively. The transition rates depend on the state of the calorimeter and are of the form

$$\Gamma_{\uparrow,k}(n) = g_k^2 \langle \Psi_n | d_k^\dagger d_k | \Psi_n \rangle, \quad (6)$$

$$\Gamma_{\downarrow,k}(n) = g_k^2 \langle \Psi_n | d_k d_k^\dagger | \Psi_n \rangle, \quad (7)$$

with $g_k^2 = \frac{2\pi}{\hbar} \kappa_k^2 \delta(\omega_0 - \epsilon_k/\hbar)$. From the point of view of the degrees of freedom of the qubit, Eq. (5) is non-Markovian as the evolution depends on both the qubit's state and the state of the calorimeter. For the total density matrix, Eq. (5) gives a master equation of the form

$$\begin{aligned} \dot{\rho}(t) &= \mathcal{L}[\rho(t)] = \frac{i}{\hbar} [\rho(t), H_q(t)] \\ &\quad - \sum_k \left\{ \frac{g_k^2}{2} [\rho(t) a a^\dagger d_k^\dagger d_k + a a^\dagger d_k^\dagger d_k \rho(t)] \right. \\ &\quad - g_k^2 a d_k^\dagger \rho(t) a^\dagger d_k - g_k^2 a^\dagger d_k \rho(t) a d_k^\dagger \\ &\quad \left. + \frac{g_k^2}{2} [\rho(t) a^\dagger a d_k d_k^\dagger + a^\dagger a d_k d_k^\dagger \rho(t)] \right\}. \end{aligned} \quad (8)$$

The results above were derived assuming that the calorimeter state can only change through the interaction with the qubit. In electronic systems [41–43], however, the internal relaxation of the calorimeter is commonly the fastest time scale. In these instances, the state of the calorimeter is more accurately described using a microcanonical ensemble instead of a single microstate. The total density matrix then reads

$$\rho(t) = \sum_E \sigma(E,t) \otimes \sigma_c(E), \quad (9)$$

where E denotes the calorimeter energy and $\sigma(E,t)$ is the qubit matrix. According to the microcanonical ensemble, the calorimeter matrix $\sigma_c(E) = [1/N(E)] \sum_k |\Psi_k\rangle\langle\Psi_k| \delta_{E_k, E}$, where E_k is the energy of microstate $|\Psi_k\rangle$ and $N(E)$ is the number of microstates with energy E . The derivation of the master equation for $\sigma(E,t)$ is similar to the previous case, yielding

$$\begin{aligned} \dot{\sigma}(E,t) &= \frac{i}{\hbar} [\sigma(E,t), H_q(t)] \\ &\quad - \left\{ \frac{\Gamma_{\uparrow}(E)}{2} [\sigma(E,t) a a^\dagger + a a^\dagger \sigma(E,t)] \right. \\ &\quad \left. - \Gamma_{\downarrow}(E - \hbar\omega_0) a \sigma(E - \hbar\omega_0, t) a^\dagger \right\} \end{aligned}$$

$$- \Gamma_{\uparrow}(E + \hbar\omega_0)a^{\dagger}\sigma(E + \hbar\omega_0,t)a + \frac{\Gamma_{\downarrow}(E)}{2}[\sigma(E,t)a^{\dagger}a + a^{\dagger}a\sigma(E,t)]\}, \quad (10)$$

where the transition rates are energy dependent $\Gamma_{\uparrow/\downarrow}(E) = [1/N(E)] \sum_{k,n} \Gamma_{\uparrow/\downarrow,k}(n)\delta_{E_n,E}$.

III. EQUIVALENCE TO THE FEQJ MODEL

In this section, we show that the same master equation (5) is also produced by the FEQJ model by averaging over the stochastic trajectories. In the FEQJ model, the interaction between the qubit and the calorimeter is described by stochastic jumps. When a jump occurs, both the qubit and calorimeter states change such that the energy difference of the calorimeter states corresponds to the energy change in the qubit. For the system studied, these jumps are caused by jump operators $D_{\downarrow,k} = g_k a \otimes d_k^{\dagger}$ and $D_{\uparrow,k} = g_k a^{\dagger} \otimes d_k$. For convenience, we denote the jump operators using only one index D_m .

Let us assume that the calorimeter is in a microstate $|\Psi_k\rangle$ and the qubit is in a state $|\psi\rangle$ at time t . According to the FEQJ protocol [36], the probability for a jump in the time interval $[t, t + \delta t]$ is given by

$$\delta p = \sum_m \delta p_m = \sum_m \delta t \text{Tr}_{q+c} \{D_m^{\dagger} D_m \sigma \otimes \sigma_c\}, \quad (11)$$

where δp_m is the probability of a jump corresponding to the jump operator D_m and the system and calorimeter states are presented in matrix form, i.e., $\sigma(t) = |\psi\rangle\langle\psi|$ and $\sigma_c(t) = |\Psi_k\rangle\langle\Psi_k|$, respectively. In the case that a jump caused by D_m occurs, the qubit state changes to $\sigma(t + \delta t) = \text{Tr}_c \{D_m \sigma \otimes \sigma_c D_m^{\dagger}\} / (\delta p_m / \delta t)$. Similarly, the calorimeter state changes to $\sigma_c(t + \delta t) = \text{Tr}_q \{D_m \sigma \otimes \sigma_c D_m^{\dagger}\} / (\delta p_m / \delta t)$.

In the case that there are no jumps in the time interval $[t, t + \delta t]$, the time evolution is given by the nonunitary Hamiltonian

$$H(t) = H_q(t) + H_c - \frac{i\hbar}{2} \sum_m D_m^{\dagger} D_m, \quad (12)$$

where H_q and H_c are the qubit and calorimeter Hamiltonians, respectively. Consequently, the qubit and calorimeter states evolve into $\sigma(t + \delta t) = \text{Tr}_c \{U(t + \delta t, t) \sigma \otimes \sigma_c U^{\dagger}(t + \delta t, t)\} / (1 - \delta p) + O(\delta t^2)$ and $\sigma_c(t + \delta t) = \text{Tr}_q \{U(t + \delta t, t) \sigma \otimes \sigma_c U^{\dagger}(t + \delta t, t)\} / (1 - \delta p) + O(\delta t^2)$, where $U(t + \delta t, t) = 1 - \frac{i}{\hbar} H(t) \delta t$. Due to the assumption that the calorimeter is in a microstate at time t , the nonunitary evolution does not change the calorimeter state, i.e., $\sigma_c(t + \delta t) = \sigma_c(t)$.

For time $t + \delta t$, we consider the total state averaged over the different outcomes between $[t, t + \delta t]$. Let us denote this averaged total state by $\rho_{\text{av}}(t + \delta t) = \sum_n \sigma_{\text{av}}(n, t + \delta t) \otimes |\Psi_n\rangle\langle\Psi_n|$, where $\sigma_{\text{av}}(n, t + \delta t)$ is the conditional average over all qubit state outcomes with calorimeter state $|\Psi_n\rangle$ multiplied with the probability of calorimeter state $|\Psi_n\rangle$. As we assume the qubit and calorimeter to be, respectively, in states $|\psi\rangle$ and $|\Psi_k\rangle$ at time t , $\sigma_{\text{av}}(n, t) = 0$ when $n \neq k$ and $\sigma_{\text{av}}(k, t) = \sigma(t) = |\psi\rangle\langle\psi|$. According to the protocol described

above,

$$\begin{aligned} \sigma_{\text{av}}(k, t + \delta t) &= \text{Tr}_c \{U(t + \delta t, t) \sigma_{\text{av}}(k, t) \otimes |\Psi_k\rangle\langle\Psi_k| U^{\dagger}(t + \delta t, t)\} \\ &= \sigma_{\text{av}}(k, t) - \frac{i}{\hbar} \delta t [H_q(t), \sigma_{\text{av}}(k, t)] \\ &\quad - \frac{1}{2} \sum_m \delta t \text{Tr}_c \{D_m^{\dagger} D_m \sigma_{\text{av}}(k, t) \otimes |\Psi_k\rangle\langle\Psi_k|\} \\ &\quad - \frac{1}{2} \sum_m \delta t \text{Tr}_c \{\sigma_{\text{av}}(k, t) \otimes |\Psi_k\rangle\langle\Psi_k| D_m^{\dagger} D_m\} \\ &\quad + O(\delta t^2), \end{aligned} \quad (13)$$

as only the no-jump evolution contributes to $\sigma_{\text{av}}(k, t + \delta t)$. Let us use $|\Psi_{k_m}\rangle$ to denote the new calorimeter microstate if a jump caused by D_m occurred from state $|\Psi_k\rangle$. Due to the possibility of these jumps, $\sigma_{\text{av}}(k_m, t + \delta t)$ becomes nonzero:

$$\sigma_{\text{av}}(k_m, t + \delta t) = \delta t \text{Tr}_c \{D_m \sigma_{\text{av}}(k, t) \otimes |\Psi_k\rangle\langle\Psi_k| D_m^{\dagger}\}. \quad (14)$$

The above results were obtained by assuming a fixed state at time t . If we now average over all the possible values of qubit and calorimeter states at time t , we get

$$\begin{aligned} \overline{\sigma_{\text{av}}}(n, t + \delta t) &= \overline{\sigma_{\text{av}}}(n, t) - \frac{i}{\hbar} \delta t [H_q(t), \overline{\sigma_{\text{av}}}(n, t)] \\ &\quad - \frac{1}{2} \sum_m \delta t \text{Tr}_c \{D_m^{\dagger} D_m \overline{\sigma_{\text{av}}}(n, t) \otimes |\Psi_n\rangle\langle\Psi_n|\} \\ &\quad - \frac{1}{2} \sum_m \delta t \text{Tr}_c \{\overline{\sigma_{\text{av}}}(n, t) \otimes |\Psi_n\rangle\langle\Psi_n| D_m^{\dagger} D_m\} \\ &\quad + \sum_m \delta t \text{Tr}_c \{D_m \overline{\sigma_{\text{av}}}(n'_m, t) \otimes |\Psi_{n'_m}\rangle\langle\Psi_{n'_m}| D_m^{\dagger}\} + O(\delta t^2) \end{aligned} \quad (15)$$

for all n , where the overbar denotes averaging over all the possible qubit and calorimeter states at time t and the index n'_m denotes the calorimeter state $|\Psi_{n'_m}\rangle$ from which the jump operator D_m can cause a jump to the calorimeter state $|\Psi_n\rangle$. By inserting the exact form of the jump operators and defining $\overline{\sigma_{\text{av}}}(n, t) = \lim_{\delta t \rightarrow 0} [\overline{\sigma_{\text{av}}}(n, t + \delta t) - \overline{\sigma_{\text{av}}}(n, t)] / \delta t$, the equation becomes equivalent to Eq. (5):

$$\begin{aligned} \dot{\overline{\sigma_{\text{av}}}}(n, t) &= \frac{i}{\hbar} [\overline{\sigma_{\text{av}}}(n, t), H_q(t)] \\ &\quad - \sum_k \left\{ \frac{\Gamma_{\uparrow,k}(n)}{2} [\overline{\sigma_{\text{av}}}(n, t) a a^{\dagger} + a a^{\dagger} \overline{\sigma_{\text{av}}}(n, t)] \right. \\ &\quad - \Gamma_{\downarrow,k}(n'_k) a \overline{\sigma_{\text{av}}}(n'_k, t) a^{\dagger} - \Gamma_{\uparrow,k}(n''_k) a^{\dagger} \overline{\sigma_{\text{av}}}(n''_k, t) a \\ &\quad \left. + \frac{\Gamma_{\downarrow,k}(n)}{2} [\overline{\sigma_{\text{av}}}(n, t) a^{\dagger} a + a^{\dagger} a \overline{\sigma_{\text{av}}}(n, t)] \right\}, \end{aligned} \quad (16)$$

with transition rates given by Eqs. (6) and (7) and the indices n'_k and n''_k satisfying $|\Psi_{n'_k}\rangle = d_k |\Psi_n\rangle / \|d_k |\Psi_n\rangle\|$ and $|\Psi_{n''_k}\rangle = d_k^{\dagger} |\Psi_n\rangle / \|d_k^{\dagger} |\Psi_n\rangle\|$, respectively. In a similar manner, when the calorimeter reaches a microcanonical ensemble immediately

after a jump, the FEQJ model produces a master equation equivalent to Eq. (10) (see Appendix B).

IV. MOMENTS OF WORK

Due to the difference in calculating thermodynamics observables, the equivalence between the master equations is not enough to guarantee that the distributions of the observables are equivalent in both methods. In this section we show that the moments of work produced by the methods are indeed identical when using the TMP without the interaction energy. However, the power operator approach is found not to agree with the TMP results without certain assumptions of the driving. We assume here that at time $t = 0$ both the qubit and the calorimeter start from thermal equilibrium with respect to inverse temperature β such that they can be expressed as a tensor product $\rho(0) = \rho_q(0) \otimes \rho_c(0)$, which is a stationary solution of Eq. (8).

A. The FEQJ method

The work done in a single FEQJ trajectory can be obtained with a projective energy measurement for both the qubit and the calorimeter in the beginning ($t = 0$) and end of the drive ($t = \tau$). The work is then defined as the energy difference between the final and initial outcomes. That is, the qubit-calorimeter interaction is neglected in the work values.

In the case of a qubit, the additional projective energy measurements for the qubit are not necessary as the calorimeter itself acts as a measurer. The initial and final energy of the qubit can be determined from the last jump before the drive and from the first jump after the drive, respectively. The total heat exchanged during the drive is given by summing over the heat exchanges caused by the jumps in the trajectory. A jump down and a jump up in the qubit cause an $\hbar\omega_0$ heat emission to the calorimeter and an $\hbar\omega_0$ heat absorption from the calorimeter, respectively. The work of a trajectory is then obtained as the change in the internal energy of the qubit plus the total heat released to the calorimeter. This leads to a work distribution equivalent to that of the double projective measurements for both the qubit and the calorimeter [36].

As shown in Appendix C, the moment generating function of the resulting work distribution can be expressed as

$$\langle e^{i\mu W_{\text{TMP}}} \rangle = \text{Tr}_{q+c} \{ \mathcal{T}_{\leftarrow} e^{i\mu [H_{q+c}^H(\tau) - H_{q+c}^H(0)]} \rho(0) \}, \quad (17)$$

where $H_{q+c}(t) = H_q(t) + H_c$ is the Hamiltonian of the composite system without the interaction Hamiltonian and $\mathcal{T}_{\leftarrow}$ is the time ordering operator. The superscript H denotes the Heisenberg picture, such that $H_{q+c}^H(t) = H_{q+c}(t)V(t,0)$, where the superoperator $V(t,0) = \mathcal{T}_{\leftarrow} \exp[\int_0^t \mathcal{L}(t')dt']$ acts on the objects on the right side of it. The moments take the form

$$\begin{aligned} \langle W_{\text{TMP}}^n \rangle &= (-i)^n \frac{\partial^n \langle e^{i\mu W_{\text{TMP}}} \rangle}{\partial \mu^n} \Big|_{\mu=0} \\ &= \text{Tr}_{c+q} \{ \mathcal{T}_{\leftarrow} \{ [H_{q+c}^H(\tau) - H_{q+c}^H(0)]^n \} \rho(0) \} \\ &= \sum_{m=0}^n \binom{n}{m} \text{Tr}_{c+q} \{ [H_{q+c}^H(\tau)]^{n-m} [H_{q+c}^H(0)]^m \rho(0) \}. \end{aligned} \quad (18)$$

These correlation functions can be calculated directly using the master equations (8) and (10). As these equations are linear in terms of the density matrices, we can express the moments as

$$\begin{aligned} \langle W_{\text{TMP}}^n \rangle &= \sum_{m=0}^n \sum_{i,k} \binom{n}{m} (-E_k - \hbar\omega_0 \delta_{i,1})^m \\ &\quad \times \text{Tr}_{c+q} \{ [H_{q+c}^H(\tau)]^{n-m} \chi(i,k,\tau) \}, \end{aligned} \quad (19)$$

where $\chi(i,k,\tau)$ is a density matrix that evolves according to the master equation with initial value $\chi(i,k,0) = |i\rangle\langle i| \otimes |\Psi_k\rangle\langle \Psi_k|$. As shown in the previous section, both methods produce equivalent evolution for the density matrix. Thus, they also produce equivalent evolution for $\chi(i,k,\tau)$ and consequently equivalent moments of work.

B. Power operator approach

For direct master equation calculations, the moments of work can be alternatively calculated with the power operator approach, which can be derived by starting from the two measurement protocol of an isolated system whose evolution is unitary during the protocol [24]. With the power operator approach, the first two work moments for the total isolated system are given by [23,24]

$$\langle W_P \rangle = \int_0^\tau dt \text{Tr}_{c+q} \{ P^{H_u}(t) \rho(0) \}, \quad (20)$$

$$\begin{aligned} \langle W_P^2 \rangle &= 2 \int_0^\tau dt_1 \int_0^{t_1} dt_2 \\ &\quad \times \text{Re} \{ \text{Tr}_{c+q} \{ P^{H_u}(t_1) P^{H_u}(t_2) \rho(0) \} \}, \end{aligned} \quad (21)$$

where $H(t)$ is the total Hamiltonian of the composite system, $P(t) = \partial_t H(t)$, and the superscript H_u denotes the usual Heisenberg picture of unitary evolution. When the driving term only acts on the qubit degrees of freedom, $P(t) = \partial_t H_q(t)$. Equations (20) and (21) can then be calculated by replacing the unitary evolution with the one given by the master equation (8) in the finite-environment case, or with a Lindblad equation in the case of an infinite environment [24]. However, as the approximations are done in a different order as compared to the way of calculating the work with the FEQJ method, the resulting moments of work are not completely identical unless certain assumptions are made on the driving.

To illustrate this, let us start from the average work given by FEQJ method, which can also be expressed as

$$\begin{aligned} \langle W_{\text{TMP}} \rangle &= \int_0^\tau dt \text{Tr}_{c+q} \{ \partial_t [H_{q+c}^H(t) \rho(0)] \} \\ &= \int_0^\tau dt \text{Tr}_{c+q} \{ \partial_t H_q(t) \rho(t) \} \\ &\quad + \int_0^\tau dt \text{Tr}_{c+q} \{ H_{q+c}(t) \dot{\rho}(t) \}, \\ &= \langle W_P \rangle + \int_0^\tau dt \text{Tr}_{c+q} \{ H_{q+c}(t) \mathcal{L}[\rho(t)] \}, \end{aligned} \quad (22)$$

where $\mathcal{L}[\rho(t)]$ is given by Eq. (8). Clearly, the average works are equivalent for all driving times τ only if $\text{Tr}_{c+q} \{ H_{q+c}(t) \mathcal{L}[\rho(t)] \}$ is zero for all t . Inserting Eq. (8), the

difference between the average work values simplifies to

$$\langle W_{\text{TMP}} \rangle - \langle W_P \rangle = - \sum_{n,k} [\Gamma_{\downarrow,k}(n) + \Gamma_{\uparrow,k}(n)] \times \int_0^\tau dt \text{Re}\{\lambda(t) \langle 0 | \sigma(n,t) | 1 \rangle\}. \quad (23)$$

For a sinusoidal driving $\lambda(t) = \lambda_0 \sin(\omega_d t)$, the term $\text{Re}\{\lambda(t) \langle 0 | \sigma(n,t) | 1 \rangle\}$ is generally nonzero. However, in the case of resonant driving, i.e., $\omega_d = \omega_0$, the term becomes zero if the fast oscillating terms of the drive are neglected. This can be shown by changing into an interaction picture with respect to H_0 :

$$\langle W_{\text{TMP}} \rangle - \langle W_P \rangle = - \sum_{n,k} [\Gamma_{\downarrow,k}(n) + \Gamma_{\uparrow,k}(n)] \times \int_0^\tau dt \text{Re}\{\lambda_I(t) \langle 0 | \sigma_I(n,t) | 1 \rangle\}, \quad (24)$$

where the driving term $\lambda_I(t) = e^{i\omega_0 t} \lambda(t)$ and I denotes the interaction picture. As we start from thermal equilibrium, $\sigma_I(n,0)$ is real for all n . In the interaction picture, the master equation (5) is real and consequently $\sigma_I(n,t)$ stays real for all t . When the fast oscillating terms in the drive are neglected, $\lambda_I(t) = i\lambda_0/2$ becomes imaginary and consequently $\text{Re}\{\lambda_I(t) \langle 1 | \sigma_I(n,t) | 0 \rangle\} = 0$. A similar difference occurs also in the higher moments.

The difference exists also when the qubit is coupled to an infinite or memoryless environment. In this case the evolution can be described with Eq. (10) by replacing all σ of different energies with $\rho_q(t)$ and removing the energy dependence of the transition rates. As the equation is then in the Lindblad form, the standard quantum jump method can be used. For the system studied, the difference between the average work becomes

$$\langle W_{\text{TMP}} \rangle - \langle W_P \rangle = -(\Gamma_\downarrow + \Gamma_\uparrow) \int_0^\tau dt \text{Re}\{\lambda_I(t) \langle 0 | \rho_{q,I}(t) | 1 \rangle\}, \quad (25)$$

where $\rho_{q,I}(t)$ is the qubit density matrix in the interaction picture with respect to H_0 , and $\Gamma_\downarrow$ and $\Gamma_\uparrow$ are the transition rates for jump down and up, respectively. Again, for a resonant sinusoidal driving, the difference becomes zero if the fast oscillating terms of the drive are neglected.

Both definitions give the same work moments also if the qubit is driven adiabatically and the jumps occur between the instantaneous eigenstates. However, in the case of nearly adiabatic off-resonance driving [28], the definitions can lead to different work moments [44].

V. NUMERICAL RESULTS

We focus on a weakly driven qubit coupled to a finite calorimeter consisting of ten two-level systems, whose energy gap is equivalent to that of the qubit. The qubit is assumed to be coupled to each two-level system with the same coupling strength. Both the qubit and the calorimeter start from thermal equilibrium such that the initial total density matrix can be

FIG. 2. Time evolution of the qubit's excited-state population given by direct master equation calculations [Eq. (10)] for different driving frequencies ω_d . The calorimeter consists of ten two-level systems with an energy gap equivalent to that of the qubit. The drive is discretized using 10^6 equidistant time steps. The driving amplitude $\lambda_0 = 0.05\hbar\omega_0$ and the coupling strength $|g|^2 = 0.01/(n_e\hbar)$, where $n_e = 10$ is the number of two-level systems in the calorimeter.

written as

$$\begin{aligned} \rho(0) &= \rho_q(0) \otimes \rho_c(0) \\ &= \sum_{i,k} \frac{e^{-\beta(E_k + \hbar\omega_0\delta_{i,1})}}{Z_q Z_c} |i\rangle\langle i| \otimes |\Psi_k\rangle\langle \Psi_k|, \end{aligned} \quad (26)$$

where β is the inverse temperature of the ideal bath, $Z_q = \sum_i e^{-\beta\hbar\omega_0\delta_{i,1}}$ is the partition function of the qubit, and $Z_c = \sum_k e^{-\beta E_k}$ is the partition function of the calorimeter. We study three sinusoidal driving protocols: a resonant driving $\omega_d = \omega_0$ and nonresonant driving frequencies $\omega_d = 0.9\omega_0$ and $\omega_d = 1.1\omega_0$. In the simulations, we use the driving amplitude $\lambda_0 = 0.05\hbar\omega_0$. For simplicity, we focus on the case where the calorimeter relaxes to a microcanonical state instantaneously after a jump. Consequently, there are only 11 calorimeter states to consider. We use the coupling strength $|g|^2 = 0.01/(n_e\hbar)$, where $n_e = 10$ is the number of two-level systems in the calorimeter.

We calculate the time evolution of the qubit density matrix directly by evolving the master equation. The time evolution of the qubit's excited-state population is illustrated in Fig. 2 for the driving frequencies $\omega_d/\omega_0 = 0.9, 1.0$, and 1.1 . As can be seen from the figure, resonant driving pumps the qubit from the ground state to the excited state and vice versa much more efficiently than the nonresonant driving frequencies $\omega_d = 0.9\omega_0$ and $1.1\omega_0$. Due to the coupling to the calorimeter, the excited state population's oscillation caused by the drive weakens in time.

In order to compare the master equation evolution to the FEQJ results, we calculate the trace distance between the total density matrices of the methods. As the total density matrices are Hermitian, the trace distance can be calculated as

$$T(\rho^{\text{ME}}, \rho^{\text{QJ}}) = \frac{1}{2} \sum_j |\mu_j|, \quad (27)$$

FIG. 3. Maximum trace distance T between the total density matrices given by the FEQJ simulations ρ^{QJ} and direct master equation calculations ρ^{ME} as a function of the number of trajectories in the FEQJ simulations. The parameters are the same as in Fig. 2.

where ρ^{ME} denotes the total density matrix given by the master equation evolution (10), ρ^{QJ} denotes the total density matrix formed from the FEQJ trajectories, and μ_j are the eigenvalues of the matrix $\rho^{ME} - \rho^{QJ}$. Figure 3 shows the maximum trace distance between the total matrices as a function of the number of trajectories in the FEQJ simulations. For all the driving frequencies studied, the FEQJ density matrix approaches the ME density matrix by increasing the number of trajectories. This agrees with the theoretical prediction that both methods give identical evolution within the numerical accuracy.

We also investigated the work statistics produced by the methods. For all the driving frequencies, we calculate the first two moments of work with both methods by using the two-measurement protocol without the interaction energy. For the direct master equation calculations, we additionally

FIG. 4. Comparison between the average work given by the power operator approach (POA) $\langle W_P \rangle$ and the two-measurement protocol $\langle W_{TMP} \rangle$ for different driving frequencies. The POA results are given by the direct master equation calculations. The lines correspond to TMP average work values given by direct master equation calculations and the markers correspond to the ones given by the FEQJ simulations. The parameters are the same as in Fig. 2. The inset shows the average work of the power operator approach given by the direct master equation calculations.

FIG. 5. Comparison between the second moment of work given by the POA $\langle W_P^2 \rangle$ and the TMP $\langle W_{TMP}^2 \rangle$ for different driving frequencies. The POA results are given by the direct master equation calculations. The lines correspond to TMP second work moment values given by direct master equation calculations and the markers correspond to the ones given by the FEQJ simulations. The parameters are the same as in Fig. 2. The inset shows the second work moment of the power operator approach given by the direct master equation calculations.

calculate the moments using the power operator approach. The results for the average work are presented in Fig. 4. The inset shows the average work given by the power operator approach. As can be seen from the inset, the resonant driving produces much larger work values than the nonresonant driving. This is due to the fact that the resonant driving pumps the qubit from the ground state to the excited state with a much higher rate as already witnessed in Fig. 2. For resonant driving, the power operator definition of work agrees well with the two-measurement-protocol results as the contribution of the fast oscillating terms in Eq. (23) is small due to weak and periodic driving. However, the power operator definition of work significantly deviates from the two-measurement-protocol results in the case of nonresonant driving. Already in the case of a slightly-off-resonance driving of $\omega_d/\omega_0 = 1.0 \pm 0.1$, the average work from the TMP and the power operator approaches differs up to 10% with the parameters studied. As proven in Sec. IV A, the FEQJ model and the direct master equation calculations are found to agree when using the TMP definition of work regardless of the driving frequency.

For the second moment of work (Fig. 5), the TMP and power operator approach are found to agree in the case of resonant driving. For the nonresonant driving of $\omega_d/\omega_0 = 1 \pm 0.1$, the approaches differ even more than in the case of the average work, up to 20%. Regardless of the driving frequency, the second work moments of the FEQJ model and the direct master equation calculations are found to agree when using the TMP definition of work as illustrated in Fig. 5.

VI. CONCLUSION

In this article we have studied the compatibility of the FEQJ model and the corresponding master equation. We have shown theoretically that the two methods produce equivalent evolution for the total density matrix. We have additionally

shown that both methods produce equivalent work moments when the work is defined using the TMP without the interaction energy. However, the power operator definition of work can deviate from the TMP work values due to a different order of approximations. In the case of adiabatic driving or sinusoidal resonant driving, the power operator approach agrees with the TMP work values.

To illustrate the results, we have numerically studied a qubit coupled to ten two-level systems with an energy gap equivalent to the one of the qubit. We have shown that the FEQJ model and the master equation produce equivalent density matrix evolution and TMP work values within the numerical accuracy. We have also shown that with the parameters studied, the average work given by the power operator approach can differ from the TMP results by 10% due to a slightly-off-resonance sinusoidal driving $\omega_d = 1.0\omega_0 \pm 0.1\omega_0$. In the case of the second moment of work, the deviation is even larger. This highlights the importance of treating work consistently with the method that is used to describe the dynamics. In the case of FEQJ, the TMP without the interaction energy appears to be the natural choice as the model itself neglects the interaction energies.

ACKNOWLEDGMENTS

We wish to thank Jukka Pekola, Mikko Möttönen, Juha Salmilehto, Shilpi Singh, and Rebecca Schmidt for useful discussions. This work was supported in part by the Väisälä Foundation and Academy of Finland through its Centres of Excellence Programme under Projects No. 251748 and No. 284621. The numerical calculations were performed using computer resources of the Aalto University School of Science “Science-IT” project.

APPENDIX A: DERIVATION OF EQ. (5) FROM EQ. (4)

By inserting the Hamiltonians into Eq. (4), it takes the form

$$\begin{aligned} \dot{\sigma}_I(n,t) = & \frac{i}{\hbar} [\sigma_I(n,t), V_I(t)] \\ & - \frac{1}{\hbar^2} \int_{-\infty}^t dt' \sum_{k,l} \kappa_k^2 (e^{-i(\omega_0-\omega_k)t-i(\omega_0-\omega_l)t'}) \\ & \times \langle \Psi_n | [ad_k^\dagger, [ad_l^\dagger, \rho_I]] | \Psi_n \rangle \\ & + e^{-i(\omega_0-\omega_k)t+i(\omega_0-\omega_l)t'} \langle \Psi_n | [ad_k^\dagger, [a^\dagger d_l, \rho_I]] | \Psi_n \rangle \\ & + e^{i(\omega_0-\omega_k)t+i(\omega_0-\omega_l)t'} \langle \Psi_n | [a^\dagger d_k, [a^\dagger d_l, \rho_I]] | \Psi_n \rangle \\ & + e^{i(\omega_0-\omega_k)t-i(\omega_0-\omega_l)t'} \langle \Psi_n | [a^\dagger d_k, [ad_l^\dagger, \rho_I]] | \Psi_n \rangle, \end{aligned} \quad (A1)$$

where the frequency $\omega_k = \epsilon_k/\hbar$. We can simplify the expression by taking into account that the time integral $\int_{-\infty}^0 dt' e^{i\omega t'} = \pi \delta(\omega) - i\mathcal{P}(1/\omega)$, where $\mathcal{P}$ is the Cauchy principal value and the imaginary part only affects the Lamb shift. Neglecting the Lamb shift,

$$\begin{aligned} \dot{\sigma}_I(n,t) = & \frac{i}{\hbar} [\sigma_I(n,t), V_I(t)] \\ & - \frac{1}{2} \sum_k g_k^2 (\langle \Psi_n | [ad_k^\dagger, [a^\dagger d_k, \rho_I]] | \Psi_n \rangle \end{aligned}$$

$$\begin{aligned} & + \langle \Psi_n | [a^\dagger d_k, [ad_k^\dagger, \rho_I]] | \Psi_n \rangle) \\ = & \frac{i}{\hbar} [\sigma_I(n,t), V_I(t)] \\ & - \frac{1}{2} \sum_k g_k^2 \{ \nu_k(n) [\sigma_I(n,t) a a^\dagger + a a^\dagger \sigma_I(n,t)] \\ & - 2\eta_k(n'_k) a \sigma_I(n'_k, t) a^\dagger - 2\nu_k(n''_k) a^\dagger \sigma_I(n''_k, t) a \\ & + \eta_k(n) [\sigma_I(n,t) a^\dagger a + a^\dagger a \sigma_I(n,t)] \}, \end{aligned} \quad (A2)$$

where $g_k^2 = 2\pi\kappa_k^2\hbar^{-2}\delta(\omega_0 - \omega_k)$, $\nu_k(\Psi_n) = \langle \Psi_n | d_k^\dagger d_k | \Psi_n \rangle$, $\eta_k(\Psi_n) = \langle \Psi_n | d_k d_k^\dagger | \Psi_n \rangle$, and n'_k and n''_k are the calorimeter microstate indices that satisfy $|\Psi_{n'_k}\rangle = d_k |\Psi_n\rangle / \|d_k |\Psi_n\rangle\|$ and $|\Psi_{n''_k}\rangle = d_k^\dagger |\Psi_n\rangle / \|d_k^\dagger |\Psi_n\rangle\|$, respectively. The expression can be simplified by defining the up and down transition rates

$$\Gamma_{\uparrow,k}(\Psi_n) = g_k^2 \nu_k(\Psi_n), \quad (A3)$$

$$\Gamma_{\downarrow,k}(\Psi_n) = g_k^2 \eta_k(\Psi_n). \quad (A4)$$

Using these transition rates the master equation takes the same form as in Eq. (5).

APPENDIX B: EQUIVALENCE BETWEEN THE MASTER EQUATION OF EQ. (10) AND THE FEQJ MODEL WHEN THE CALORIMETER RELAXES INSTANTANEOUSLY TO A MICROCANONICAL STATE AFTER A JUMP

Let us assume that the internal relaxation of the calorimeter is the fastest time scale such that the calorimeter relaxes instantaneously after a jump into a microcanonical ensemble of the states corresponding to the same energy and stays in the microcanonical ensemble until another jump. We can still use the FEQJ model to calculate the qubit dynamics by using an averaged calorimeter state instead of a single calorimeter microstate in Eqs. (11) and (12).

Let us assume that the calorimeter has energy E at time t . For time $t + \delta t$, where δt is very short, we consider again the averaged total state $\rho_{\text{av}}(t + \delta t) = \sum_n \sigma_{\text{av}}(E_n, t + \delta t) \otimes \sigma_c(E_n)$, where $\sigma_{\text{av}}(E_n, t + \delta t)$ is the conditional average over all qubit state outcomes with calorimeter energy E_n multiplied with the probability of calorimeter energy E_n . As we assume the calorimeter to have an energy E at time t , $\sigma_{\text{av}}(E_n, t) = 0$ when $E_n \neq E$. According to the FEQJ protocol [36],

$$\begin{aligned} \sigma_{\text{av}}(E, t + \delta t) &= \text{Tr}_c \{ U(t + \delta t, t) \sigma_{\text{av}}(E, t) \otimes \sigma_c(E) U^\dagger(t + \delta t, t) \} \\ &= \sigma_{\text{av}}(E, t) - \frac{i}{\hbar} \delta t [H_q(t), \sigma_{\text{av}}(E, t)] \\ &\quad - \frac{1}{2} \sum_m \sum_{k \in \{\uparrow, \downarrow\}} \delta t \text{Tr}_c \{ D_{m,k}^\dagger D_{m,k} \sigma_{\text{av}}(E, t) \otimes \sigma_c(E) \} \\ &\quad - \frac{1}{2} \sum_m \sum_{k \in \{\uparrow, \downarrow\}} \delta t \text{Tr}_c \{ \sigma_{\text{av}}(E, t) \otimes \sigma_c(E) D_{m,k}^\dagger D_{m,k} \} \\ &\quad + O(\delta t^2), \end{aligned} \quad (B1)$$

as only the no-jump evolution contributes to $\sigma_{\text{av}}(E, t + \delta t)$. Due to the possibility of a jump corresponding to energy

change $\hbar\omega_0$, $\sigma_{\text{av}}(E + \hbar\omega_0, t + \delta t)$ and $\sigma_{\text{av}}(E - \hbar\omega_0, t + \delta t)$ become nonzero:

$$\sigma_{\text{av}}(E + \hbar\omega_0, t + \delta t) = \delta t \text{Tr}_c \left\{ \sum_m D_{m,\downarrow} \sigma_{\text{av}}(E, t) \otimes \sigma_c(E) D_{m,\downarrow}^\dagger \right\}, \quad (\text{B2})$$

$$\sigma_{\text{av}}(E - \hbar\omega_0, t + \delta t) = \delta t \text{Tr}_c \left\{ \sum_m D_{m,\uparrow} \sigma_{\text{av}}(E, t) \otimes \sigma_c(E) D_{m,\uparrow}^\dagger \right\}. \quad (\text{B3})$$

Averaging over all the trajectories with different qubit and calorimeter states at time t , we get

$$\begin{aligned} \overline{\sigma_{\text{av}}}(E_n, t + \delta t) &= \overline{\sigma_{\text{av}}}(E_n, t) - \frac{i}{\hbar} \delta t [H_q(t), \overline{\sigma_{\text{av}}}(E_n, t)] \\ &\quad - \frac{1}{2} \sum_m \sum_{k \in \{\uparrow, \downarrow\}} \delta t \text{Tr}_c \{ D_{m,k}^\dagger D_{m,k} \overline{\sigma_{\text{av}}}(E_n, t) \otimes \sigma_c(E_n) \} \\ &\quad - \frac{1}{2} \sum_m \sum_{k \in \{\uparrow, \downarrow\}} \delta t \text{Tr}_c \{ \overline{\sigma_{\text{av}}}(E_n, t) \otimes \sigma_c(E_n) D_{m,k}^\dagger D_{m,k} \} \\ &\quad + \sum_m \delta t \text{Tr}_c \{ D_{m,\downarrow} \overline{\sigma_{\text{av}}}(E_n - \hbar\omega_0, t) \otimes \sigma_c(E_n - \hbar\omega_0) D_{m,\downarrow}^\dagger \} \\ &\quad + \sum_m \delta t \text{Tr}_c \{ D_{m,\uparrow} \overline{\sigma_{\text{av}}}(E_n + \hbar\omega_0, t) \otimes \sigma_c(E_n + \hbar\omega_0) D_{m,\uparrow}^\dagger \} \\ &\quad + O(\delta t^2). \end{aligned} \quad (\text{B4})$$

By inserting the exact form of the jump operators and defining $\overline{\sigma_{\text{av}}}(k, t) = \lim_{\delta t \rightarrow 0} [\overline{\sigma_{\text{av}}}(k, t + \delta t) - \overline{\sigma_{\text{av}}}(k, t)] / \delta t$, the equation becomes equivalent to Eq. (5):

$$\begin{aligned} \dot{\overline{\sigma_{\text{av}}}}(E_n, t) &= \frac{i}{\hbar} [\overline{\sigma_{\text{av}}}(E_n, t), H_q(t)] \\ &\quad - \left\{ \frac{\Gamma_\uparrow(E)}{2} [\overline{\sigma_{\text{av}}}(E_n, t) a a^\dagger + a a^\dagger \overline{\sigma_{\text{av}}}(E_n, t)] \right. \\ &\quad - \Gamma_\downarrow(E_n - \hbar\omega_0) a \overline{\sigma_{\text{av}}}(E_n - \hbar\omega_0, t) a^\dagger \\ &\quad - \Gamma_\uparrow(E_n + \hbar\omega_0) a^\dagger \overline{\sigma_{\text{av}}}(E_n + \hbar\omega_0, t) a \\ &\quad \left. + \frac{\Gamma_\downarrow(E_n)}{2} [\overline{\sigma_{\text{av}}}(E_n, t) a^\dagger a + a^\dagger a \overline{\sigma_{\text{av}}}(E_n, t)] \right\}, \end{aligned} \quad (\text{B5})$$

with transition rates given by $\Gamma_{\uparrow/\downarrow}(E) = [1/N(E)] \sum_{k,n} \Gamma_{\uparrow/\downarrow,k}(n) \delta_{E_n, E}$.

APPENDIX C: WORK AMOUNTS GIVEN BY THE FEQJ METHOD

For simplicity, let us focus on the case where the calorimeter stays in the same microstate between the jumps. We denote the probability of a quantum trajectory with N jumps by $P_{\text{QJ}}[i, f, \Psi_0, \Psi_N, \{D_{m_k}\}_{k=1}^N, \{t_k\}_{k=1}^N]$, where i is the initial state of the qubit, f is the final state of the qubit, Ψ_0 is the initial state of the calorimeter, Ψ_N is the final state of the calorimeter, and a jump caused by $D_{m_k} = g_{m_k} A_{m_k} \otimes B_{m_k}$ occurs at time t_k . As the calorimeter's state does not change between the jumps we can use the calorimeter traced jump operators $C_m = \sqrt{\Gamma_m(\Psi)} A_m$,

where Ψ is the calorimeter state and the transition rate $\Gamma_m(\Psi)$ is defined such that $\text{Tr}_q \{ C_m \sigma C_m^\dagger \} = \text{Tr}_{q+c} \{ D_m \sigma \otimes |\Psi\rangle\langle\Psi| D_m^\dagger \}$ for all qubit states $\sigma = |\psi\rangle\langle\psi|$.

Using the calorimeter traced jump operators, the trajectory's probability can be written as [36]

$$\begin{aligned} P_{\text{QJ}}[i, f, \Psi_0, \Psi_N, \{D_{m_k}\}_{k=1}^N, \{t_k\}_{k=1}^N] &= (\delta t)^N P[i, \Psi_0] \left[\prod_{k=1}^N \Gamma_{m_k}(\Psi_{k-1}) \right] \left| \langle f | U_{\text{eff}}(\tau, t_N) \right. \\ &\quad \left. \times \left[\prod_{k=1}^N A_{m_{N+1-k}} U_{\text{eff}}(t_{N+1-k}, t_{N-k}) \right] | i \rangle \right|^2, \end{aligned} \quad (\text{C1})$$

where $|\Psi_k\rangle$ is the calorimeter state after the k th jump and the no-jump evolution is given by

$$U_{\text{eff}}(t_{k+1}, t_k) = \mathcal{T}_\leftarrow \exp \left[-\frac{i}{\hbar} \int_{t_k}^{t_{k+1}} \left(H_q(t) - i \frac{\hbar}{2} \sum_i \Gamma_i(\Psi_k) A_i^\dagger A_i \right) dt \right],$$

where $H_q(t)$ is the qubit Hamiltonian and $\mathcal{T}_\leftarrow$ is the time ordering operator. The TMP moments of work are then given by [36]

$$\begin{aligned} \langle W_{\text{TMP}}^n \rangle &= \sum_{\text{traj}} [\hbar\omega_f + E_N - (\hbar\omega_i + E_0)]^n \\ &\quad \times P_{\text{QJ}}[i, f, \Psi_0, \Psi_N, \{D_{m_k}\}_{k=1}^N, \{t_k\}_{k=1}^N] \\ &= \sum_{\text{traj}} \sum_{k=0}^n \binom{n}{k} (\hbar\omega_f + E_N)^{n-k} (-\hbar\omega_i - E_0)^k \\ &\quad \times P_{\text{QJ}}[i, f, \Psi_0, \Psi_N, \{D_{m_k}\}_{k=1}^N, \{t_k\}_{k=1}^N]. \end{aligned} \quad (\text{C2})$$

By summing over the trajectories that start from the same initial states and produce equivalent final states, we can express the moments as

$$\begin{aligned} \langle W_{\text{TMP}}^n \rangle &= \sum_{k=0}^n \sum_{i, f, \Psi_0, \Psi_N} \binom{n}{k} (\hbar\omega_f + E_N)^{n-k} (-\hbar\omega_i - E_0)^k \\ &\quad \times \text{Tr}_{c+q} \{ |f\rangle\langle f| \otimes |\Psi_N\rangle\langle\Psi_N| \chi(i, \Psi_0, \tau) \} \\ &= \sum_{k=0}^n \sum_{i, \Psi_0} \binom{n}{k} (-\hbar\omega_i - E_0)^k \\ &\quad \times \text{Tr}_{c+q} \{ H_{q+c}^{n-k}(\tau) \chi(i, \Psi_0, \tau) \}, \end{aligned} \quad (\text{C3})$$

where $H_{q+c}(t) = H_q(t) + H_c$ is the Hamiltonian of the composite system without the interaction Hamiltonian and $\chi(i, \Psi_0, \tau)$ is a density matrix of the total system evolved according to the master equation (8) with initial value $\chi(i, \Psi_0, 0) = |i\rangle\langle i| \otimes |\Psi_0\rangle\langle\Psi_0|$. As Eq. (8) is linear, we can simplify the expression to be

$$\langle W_{\text{TMP}}^n \rangle = \sum_{k=0}^n \binom{n}{k} \text{Tr}_{c+q} \{ H_{q+c}^{n-k}(\tau) V(\tau, 0) H_{q+c}^k(0) \rho(0) \}, \quad (\text{C4})$$

where $\rho(0)$ is the initial total density matrix operated by $H_{q+c}^k(0)$ at time $t = 0$. The resulting matrix is evolved in

time according to Eq. (8). This time evolution is given by the superoperator $V(t,0) = \mathcal{T}_{\leftarrow} \exp[\int_0^t \mathcal{L}(t')dt']$ that acts on the objects on the right side of it. The corresponding moment generating function is given by

$$\langle e^{i\mu W_{\text{TMP}}} \rangle = \text{Tr}_{q+c} \left\{ \mathcal{T}_{\leftarrow} e^{i\mu [H_{q+c}^H(\tau) - H_{q+c}^H(0)]} \rho(0) \right\}, \quad (\text{C5})$$

where the superscript H denotes the Heisenberg picture such that $H_{q+c}^H(t) = H_{q+c}(t)V(t,0)$. In the case that the calorimeter instantaneously reaches a microcanonical ensemble after a jump, the derivation is similar and leads to the same form of the moment generating function.

-
- [1] M. Esposito and S. Mukamel, *Phys. Rev. E* **73**, 046129 (2006).
 - [2] M. Campisi, P. Talkner, and P. Hänggi, *Phys. Rev. Lett.* **102**, 210401 (2009).
 - [3] G. E. Crooks, *J. Stat. Mech.* (2008) P10023.
 - [4] R. Chetrite and K. Mallick, *J. Stat. Phys.* **148**, 480 (2012).
 - [5] T. Albash, D. A. Lidar, M. Marvian, and P. Zanardi, *Phys. Rev. E* **88**, 032146 (2013).
 - [6] M. Esposito, U. Harbola, and S. Mukamel, *Rev. Mod. Phys.* **81**, 1665 (2009).
 - [7] M. Campisi, P. Hänggi, and P. Talkner, *Rev. Mod. Phys.* **83**, 771 (2011).
 - [8] A. E. Rastegin and K. Życzkowski, *Phys. Rev. E* **89**, 012127 (2014).
 - [9] G. Watanabe, B. P. Venkatesh, P. Talkner, M. Campisi, and P. Hänggi, *Phys. Rev. E* **89**, 032114 (2014).
 - [10] J. Salmilehto, P. Solinas, and M. Möttönen, *Phys. Rev. E* **89**, 052128 (2014).
 - [11] J. Ankerhold and J. P. Pekola, *Phys. Rev. B* **90**, 075421 (2014).
 - [12] C. Jarzynski, H. T. Quan, and S. Rahav, *Phys. Rev. X* **5**, 031038 (2015).
 - [13] T. B. Batalhão, A. M. Souza, L. Mazzola, R. Auccaise, R. S. Sarthour, I. S. Oliveira, J. Goold, G. De Chiara, M. Paternostro, and R. M. Serra, *Phys. Rev. Lett.* **113**, 140601 (2014).
 - [14] S. An, J.-N. Zhang, M. Um, D. Lv, Y. Lu, J. Zhang, Z.-Q. Yin, H. T. Quan, and K. Kim, *Nat. Phys.* **11**, 193 (2015).
 - [15] M. Carrega, P. Solinas, A. Braggio, M. Sassetti, and U. Weiss, *New J. Phys.* **17**, 045030 (2015).
 - [16] R. Schmidt, M. F. Carusela, J. P. Pekola, S. Suomela, and J. Ankerhold, *Phys. Rev. B* **91**, 224303 (2015).
 - [17] B. P. Venkatesh, G. Watanabe, and P. Talkner, *New J. Phys.* **17**, 075018 (2015).
 - [18] J. M. Horowitz, *Phys. Rev. E* **85**, 031110 (2012).
 - [19] F. W. J. Hekking and J. P. Pekola, *Phys. Rev. Lett.* **111**, 093602 (2013).
 - [20] J. M. Horowitz and J. M. R. Parrondo, *New J. Phys.* **15**, 085028 (2013).
 - [21] B. Leggio, A. Napoli, A. Messina, and H.-P. Breuer, *Phys. Rev. A* **88**, 042111 (2013).
 - [22] F. Liu, *Phys. Rev. E* **89**, 042122 (2014).
 - [23] P. Solinas, D. V. Averin, and J. P. Pekola, *Phys. Rev. B* **87**, 060508 (2013).
 - [24] S. Suomela, P. Solinas, J. P. Pekola, J. Ankerhold, and T. Ala-Nissila, *Phys. Rev. B* **90**, 094304 (2014).
 - [25] J. Horowitz and T. Sagawa, *J. Stat. Phys.* **156**, 55 (2014).
 - [26] F. Liu, *Phys. Rev. E* **93**, 012127 (2016).
 - [27] J. P. Pekola, Y. Masuyama, Y. Nakamura, J. Bergli, and Y. M. Galperin, *Phys. Rev. E* **91**, 062109 (2015).
 - [28] S. Suomela, J. Salmilehto, I. G. Savenko, T. Ala-Nissila, and M. Möttönen, *Phys. Rev. E* **91**, 022126 (2015).
 - [29] G. Manzano, J. M. Horowitz, and J. M. R. Parrondo, *Phys. Rev. E* **92**, 032129 (2015).
 - [30] P. Talkner and P. Hänggi, *Phys. Rev. E* **93**, 022131 (2016).
 - [31] M. Carrega, P. Solinas, M. Sassetti, and U. Weiss, *Phys. Rev. Lett.* **116**, 240403 (2016).
 - [32] Z. Gong, Y. Ashida, and M. Ueda, *Phys. Rev. A* **94**, 012107 (2016).
 - [33] C. Elouard, D. H. Marti, M. Clusel, and A. Auffèves, *arXiv:1603.07266*.
 - [34] F. Jin, R. Steinigeweg, H. De Raedt, K. Michielsen, M. Campisi, and J. Gemmer, *Phys. Rev. E* **94**, 012125 (2016).
 - [35] J. P. Pekola, P. Solinas, A. Shnirman, and D. V. Averin, *New J. Phys.* **15**, 115006 (2013).
 - [36] S. Suomela, A. Kutvonen, and T. Ala-Nissila, *Phys. Rev. E* **93**, 062106 (2016).
 - [37] J. Kurchan, *arXiv:cond-mat/0007360*.
 - [38] H. Tasaki, *arXiv:cond-mat/0009244*.
 - [39] S. Mukamel, *Phys. Rev. Lett.* **90**, 170604 (2003).
 - [40] W. H. Zurek, *Rev. Mod. Phys.* **75**, 715 (2003).
 - [41] A. Kutvonen, T. Ala-Nissila, and J. Pekola, *Phys. Rev. E* **92**, 012107 (2015).
 - [42] J. Pekola, S. Suomela, and Y. Galperin, *J. Low Temp. Phys.* **184**, 1015 (2016).
 - [43] J. P. Pekola, P. Muratore-Ginanneschi, A. Kupiainen, and Y. M. Galperin, *Phys. Rev. E* **94**, 022123 (2016).
 - [44] S. Suomela, J. Salmilehto, I. G. Savenko, T. Ala-Nissila, and M. Möttönen (unpublished).